

Age-Related Changes in the Left Ventricular Twist

Sabiye Yılmaz¹, Harun Kılıç², Saadet Demirtaş¹, Mehmet Bülent Vatan², Efe Edem¹, Mustafa Tarık Ağaç², Mustafa Türker Pabuccu¹, Mehmet Akif Çakar², Mustafa Gökhan Vural², Murat Aksoy², Ersan Tatlı², Hüseyin Gündüz², Ramazan Akdemir²

1 Sakarya Eğitim ve Araştırma Hastanesi, Kardiyoloji Kliniği, Samsun

2 Sakarya Üniversitesi Tıp Fakültesi, Kardiyoloji Anabilim Dalı, Samsun

ABSTRACT

Introduction: Left ventricular (LV) rotation and twist play an important role in LV contraction and relaxation. Systolic function is usually preserved with aging but diastolic function usually deteriorates. The aim of this study was to establish a reference value for the LV twist and examine the effect of aging on twist.

Materials and Method: We enrolled 75 healthy subjects who were divided into two groups according to age: young < 40 years and older >40. The LV rotation and twist were assessed by 2D-STI at the basal and apical levels of the parasternal short-axis. LV twist was defined as an apical rotation relative to the baseline. Exclusive criteria included a history of ischemic and valvular disease, arrhythmia, a pacemaker and systolic dysfunction.

Results: Peak A wave velocity (70.6 ± 13.7 vs 57.7 ± 10.6 , $p < 0.001$), E/E' ratio (7.7 ± 2.2 vs 6.1 ± 1.3 , $p < 0.001$) were higher and peak E wave velocity (52.4 ± 16 vs 76.8 ± 11 , $p < 0.001$), E/A ratio (0.75 vs 1.34 , $p < 0.001$) were lower in old patients compared to young. There was an increase in apical rotation ($4.4 \pm 2.9^\circ$ vs $3.3 \pm 2.4^\circ$, $p = 0.075$) and a decrease in basal rotation ($-4.1 \pm 2.7^\circ$ vs $-5.2 \pm 3.7^\circ$, $p = 0.185$) in old patients, but these changes didn't indicate significant difference. With aging, the twist increased ($8.6 \pm 3.3^\circ$ vs $8.5 \pm 4.1^\circ$, $p = 0.890$) insignificantly. The LV apical rotation (5 ± 3.2 vs 2.4 ± 1.1 , $p = 0.004$) and twist (9.4 ± 3.4 vs 6.9 ± 2.4 , $p = 0.64$) decreased in the old group with an increasing degree of LV diastolic dysfunction.

Conclusion: LV twist increased slightly with age, but this effect was reversed with increased diastolic dysfunction even in the presence of normal systolic function.

Keywords: Echocardiography, rotation and twist, aging

Sol Ventrikül Bükülme Hareketinde Yaşa Bağlı Değişiklikler

Sabiye Yılmaz¹, Harun Kılıç², Saadet Demirtaş¹, Mehmet Bülent Vatan², Efe Edem¹, Mustafa Tarık Ağaç², Mustafa Türker Pabuccu¹, Mehmet Akif Çakar², Mustafa Gökhan Vural², Murat Aksoy², Ersan Tatlı², Hüseyin Gündüz², Ramazan Akdemir²

1 Sakarya Eğitim ve Araştırma Hastanesi, Kardiyoloji Kliniği, Samsun

2 Sakarya Üniversitesi Tıp Fakültesi, Kardiyoloji Anabilim Dalı, Samsun

ÖZET

Giriş: Sol ventrikül (LV) rotasyon ve büküm hareketi kasılma ve gevşemede önemli rol oynamaktadır. Yaşlanma ile LV sistolik fonksiyonları korunsada diyastolik fonksiyonlar genellikle bozulur. Bu çalışmanın amacı, LV büküm için bir referans değeri saptamak ve yaşlanmanın büküm üzerinde etkilerini incelemektir.

Hastalar ve Metod: Çalışmaya 75 sağlıklı kişi dahil edildi; katılımcılar 40 yaş üstü ve 40 yaş altı olarak iki gruba ayrıldı. LV rotasyon ve büküm parasternal kısa eksen bazal ve apikal seviyelerinde 2D-STI ile değerlendirildi. LV büküm hareketi apikal ve bazal rotasyonun göreceli toplamı olarak tanımlandı. İskemik ve kapak hastalığı, aritmi, kalp pili ve sistolik disfonksiyon öyküsü dışlama kriterleri olarak tanımlandı.

Bulgular: Yaşlılarda gençlere kıyasla pik A dalga hızı (70.6 ± 13.7 vs 57.7 ± 10.6 , $p < 0.001$), E/E' oranı (7.7 ± 2.23 vs 6.09 ± 1.3 , $p < 0.001$) daha yüksek, pik E dalga hızı (52.4 ± 16 vs 76.75 ± 11 , $p < 0.001$), E/A oranı (0.75 vs 1.34 , $p < 0.001$) daha düşüktü. Yaşlı grupta apikal rotasyonda artış (4.4 ± 2.9 ° vs. 3.3 ± 2.4 °, $p = 0.075$) ve bazal rotasyonda azalma (-4.1 ± 2.7 ° vs 3.7 ± 5.2 , $p = 0.185$) izlendi ancak, bu değişiklikler anlamlı değildi. Yaşlanma ile birlikte, büküm önemsiz derecede arttı (8.6 ± 3.3 ° vs 8.53 ± 4.1 °, $p = 0.890$). Yaşlı grupta LV diyastolik disfonksiyon derecesi arttıkça apikal rotasyon (5.02 ± 3.2 vs 2.4 ± 1.1 , $p = 0.004$) ve büküm (9.4 ± 3.4 vs 6.9 ± 2.4 , $p = 0.64$) azaldı.

Sonuç: Bu çalışma LV bükümün yaşla birlikte hafifçe arttığını, ancak bu etkinin sistolik fonksiyonları normal olsada artmış diyastolik disfonksiyon varlığında tersine döndüğünü gösterdi.

Anahtar Kelimeler: Ekokardiyografi, rotasyon ve büküm, yaşlanma

Geliş Tarihi: 09.06.2016 - **Kabul Tarihi:** 21.09.2016

INTRODUCTION

The LV has an oblique myocardial fibre orientation that induces rotation around the long axis during contraction [1]. The LV twist is defined as a wringing motion of the heart caused by the opposite rotation of the LV apical and basal segments [1,2]. The LV rotation and twist are essential components of cardiac performance [3]. Measurement of the LV twist may play an important role in determining the LV systolic and diastolic dysfunction [1,4]. Alterations in the LV twist have been reported in several pathophysiological conditions, and the potential clinical applications of the twist are currently expanding [5–8]. LV systolic function is usually preserved with aging, but diastolic function usually deteriorates. Evaluation of the LV rotation and twist are feasible with two-dimensional speckle tracking imaging (2D-STI) [9-10].

The aim of this study was to establish a reference value of LV twist and to examine the effect of aging on LV twist in a group of healthy volunteers.

METHODS

Patient selection

Initially, the study population consisted of 102 healthy volunteers, but 27 were excluded because of insufficient image quality, and the remaining 75 subjects (62%) made up the final study group. The study population was divided into two age groups: < 40 years (n: 40) and >40 years (n:35). A history of heart disease, such as ischemic and valvular disease, arrhythmia, pacemaker or systolic dysfunction, was considered an exclusion factor. Seventeen older patients were receiving antihypertensive therapy, and their blood pressure was under control. They had normal electrocardiographic findings.

The study was approved by the local institutional ethics committee.

Echocardiography

A standard comprehensive M-mode, 2D echocardiogram, Doppler study were performed with the Philips IE33 system (Philips Medical Systems, Amsterdam, Netherlands) using a S5-1 transducer. The peak early (E) and late (A) filling velocities and the E/A ratio were measured using pulsed-wave Doppler of the mitral inflow. Mitral annulus velocities were achieved from the septal and lateral annulus of the left ventricle by tissue Doppler imaging (TDI).The patients were stratified by the grade of

diastolic dysfunction that was classified as having impaired relaxation (grade 1), pseudonormal (grade 2) or restrictive patterns of echocardiographic diastolic filling. In addition, all patients with diastolic dysfunction were required to have early diastolic velocity of the septal mitral annulus (E') of less than 8 cm/s [11].

After a comprehensive standard echocardiographic examination, apical and basal short-axis rotations were measured by 2D-STI. Short axis basal image was defined from the mitral ring level, and the apical image was defined by the LV cavity alone with no papillary muscle visible [12-13]. At each plane, four sequential cardiac cycles were obtained and digitally stored in a hard disc for offline analysis. Using commercially available 2D strain software, the endocardial border of each short axis plane in the end-systolic frame was manually traced. The software algorithm then automatically segmented the LV short-axis plane into six segments and searched speckles in the ROIs on a frame-by-frame basis using the sum of absolute difference algorithm. Afterwards, software defined the ventricular centroid for the mid-myocardial line on a frame-by-frame basis during one cardiac cycle, and it calculated the time domain LV rotation and radial displacement profiles for each segment in both short axis planes [14,15]. LV twist was calculated as the apical LV rotation relative to the basal rotation.

Statistical Analysis

Two independent sample t-tests were used to compare the continuous data between the two groups according to the parametrical assumed. Continuous data were presented as the mean \pm standard. Categorical data were compared by chi-square tests. Categorical variables were presented as a count and percentage. Pearson's correlation coefficients was used for the determination of the correlations between the LV twist and other parameters. A p-value <0.05 was considered significant. Analyses were performed using commercial software (IBM Corp. IBM SPSS Statistics for Windows, Version 16.0. Armonk, NY).

RESULTS

The clinical characteristics and echocardiographic parameters of the two age groups are shown in Table 1. Although left ventricular ejection fraction (LVEF) was preserved in all participants, LVEF was higher in the young group ($p=0.045$). The rate of diabetes mellitus ($p=0.03$), obesity ($p: 0.01$) and

hypertension ($p < 0.000$) were higher in the elderly group, but blood pressure was controlled in these patients. Variables representing the LV diastolic function were found to be distinctly different between the two groups. As expected, Doppler measurements revealed that the peak A wave velocity (70.6 ± 13.7 vs. 57.7 ± 10.6 , $p < 0.001$) and E/E' ratio (7.7 ± 2.2 vs. 6.1 ± 1.3 , $p < 0.001$) increased, and the peak E wave velocity (52.4 ± 16 vs. 76.8 ± 11 , $p < 0.001$), early diastolic mitral annular velocity (E' wave) (7 ± 1.9 vs. 13.1 ± 3.6 , $p = 0.00$) and E/A ratio (0.75 vs. 1.3 , $p < 0.001$) decreased significantly with advancing aging. Compared with the young group, there was increase in magnitude of peak systolic apical rotation ($4.4 \pm 2.9^\circ$ vs. $3.3 \pm 2.4^\circ$, $p = 0.075$) and a decrease in basal rotation ($-4.1 \pm 2.7^\circ$ vs. $-5.2 \pm 3.7^\circ$, $p = 0.185$) in the older group, but it was not significant. The peak LV twist increased with advancing aging insignificantly; the mean value of peak twist was observed $8.5 \pm 4.1^\circ$ (ranged between 4.4° and 12.6°) for < 40 year old participants and $8.7 \pm 3.3^\circ$ (ranged between 5.3° and 12.9°) for > 40 year old participants ($p = 0.890$). In the univariate analysis, LV twist was insignificantly correlated with age ($r = 0.081$, $p = 0.491$), BMI ($r = 0.047$, $p = 0.689$), diabetes mellitus ($r = 0.050$; $p = 0.672$) and hypertension ($r = 0.146$; $p = 0.212$). In addition, basal rotation, apical rotation and LV twist were not affected by LVEF. LV apical rotation (normal = $4.1 \pm 2.5^\circ$ vs. grade 1 = $5.2 \pm 3.2^\circ$, $p = 0.699$), and twist (normal = $6.9 \pm 2.4^\circ$ vs. grade 1 = $9.7 \pm 3.2^\circ$, $p = 0.084$) were increased in patients with mild diastolic dysfunction and were decreased in more advanced diastolic dysfunction (apical rotation in grade 2 was $2.4 \pm 1.2^\circ$, $p = 0.530$ vs. the twist in grade 2 which was $6.7 \pm 3.3^\circ$, $p = 0.990$) compared with normal healthy people (Table 2).

The correlation was not significant between the LV twist and the marker of increased LV end-diastolic pressure: E/A ratio ($r = 0.156$, $p = 0.180$) and E/E' ($r = 0.135$, $p = 0.246$).

DISCUSSION

This study has investigated the normal reference values of principal LV with parameters, and analyzed the age-related changes in the LV twist. LV rotation and twist were increased with age because of increases in apical rotations, but this relationship was not statistically significant. Moreover, an association between the LV twist and diastolic dysfunction was shown in this study; the peak LV twist was increased in patients with mild or early-stage diastolic dysfunction and it was normalized or decreased with an increasing degree of diastolic dysfunction. These findings may provide supplementary information regarding the effect of diastolic dysfunction on the LV twist.

Cardiac Motion and Twist

Normally, LV performs a wringing systolic motion with a clockwise rotation in the basal plane and a counterclockwise rotation in the apical plane [14,15]. As shown in Figure 1, viewed from the LV apex, counterclockwise rotation was expressed as a positive value and clockwise rotation was expressed as a negative value. Any factor that impoverished the LV subendocardial fibres can lead to an increased twist. For example, the systolic twist increases in hypertrophic cardiomyopathy or LV hypertrophy caused by aortic stenosis [16-18]. Both conditions lead to a subendocardial dysfunction, causing an increased LV twist [19]. Though the primary pathologic mechanisms for age-related twist changes are not completely known, similarly, it has been proposed that the aging process causes subendocardial fibrosis and a decline in subendocardial function [19-21]. Thus, optimization of the LV twist is an effective method to preserve the LV ejection fraction in the elderly. Our study showed that the LV rotation and twist gradually increase with age, which these results are in accordance with previous studies [19-22]. Evaluation of the LV rotation and twist is feasible with 2D-STI. However, LV twist value determined by 2D-STI has significant inter-study discrepancies. For example, Takeuchi et al. [19] and Saygisunar U et al [7] reported a twist value in the range of 6° to 7°, which is similar to the values in our study ($8.5\pm 4.1^\circ$ vs. $8.7\pm 3.3^\circ$). Lee et al. [22] and Sun et al. [20] reported peak twist values in the range of 20°, which was higher than the peak LV twist reported in our study. The reasons for this inconsistency may be due to the selection of an apical basal plane and the echocardiographic image quality resolution.

LV Twist and Diastolic Dysfunction

Our study showed that the LV twist was dependent on the degree of diastolic dysfunction. Consistent with previous study [19], the peak LV twist was increased in patients with mild or early-stage diastolic dysfunction, but it was normalized or decreased with an increasing degree of diastolic dysfunction compared with normal healthy controls. LV twist was also found to be increased in asymptomatic elderly patients whose diastolic function was generally mildly reduced compared with younger people. This suggests that an increased LV twist is a part of the inherent progression of diastolic dysfunction, and to preserve normal LV filling, it is the first compensation for the decreased function of the myocardium. The increased twist can be used as a marker for the early stage of diastolic dysfunction with normal filling pressure, but this requires more investigation.

Limitations of the study

Our study was limited by the relatively small sample size. Another limitation of this study was the poor quality of some of the 2D images. The selection of optimal imaging planes is quite challenging because of limited acoustic windows and the oblique orientation of the heart in the chest cavity. A small proportion of patients had hypertension, which may have affected the LV mechanics. The main purpose of this study was not to investigate the LV twist characteristics with different grades of diastolic dysfunction, so patients with grade 3 diastolic dysfunction were not included in this study.

CONCLUSIONS

An evaluation of LV rotation and twist is feasible with 2D-STI. We assessed the LV rotation and twist mechanics in young and older people to determine a value for the twist in a normal adult population. This value can serve as a reference for further evaluation of pathological myocardial motions in various cardiovascular and other diseases [23-25]. In addition, our data suggest that this non-invasive parameter can provide helpful information in the assessment of diastolic function. An increased twist can also be used as a marker for the early stage of diastolic dysfunction with normal filling pressure, as the twist may decrease with an increasing degree of diastolic dysfunction. We need further studies to determine the most practical and reliable parameters to identify diastolic dysfunction.

Conflict of Interests

"I declare that none of the authors have a Conflict of Interests."

REFERENCES:

1. Hollingsworth KG, Blamire AM, Keavney BD, Macgowan GA. Left ventricular torsion, energetics, and diastolic function in normal human aging. *Am J Physiol Heart Circ Physiol* 2012; 302: 885– 892.
2. Cameli M, Lisi M, Righini FM, Massoni A, Mondillo S. Left ventricular remodeling and torsion dynamics in hypertensive patients. *Int J Cardiovasc Imaging* 2013; 29: 79–86.
3. Lamia B, Tanabe M, Tanaka H, Kim HK, Gorcsan J 3rd, Pinsky MR. Left ventricular systolic torsion correlates global cardiac performance during dyssynchrony and cardiac resynchronization therapy. *Am J Physiol Heart Circ Physiol* 2011; 300: 853-8
4. Notomi Y, Srinath G, Shiota T, Martin-Miklovic MG, Beachler L, Howell K, et al. Maturation and adaptive modulation of left ventricular torsional biomechanics: Doppler tissue imaging observation from infancy to adulthood. *J Am Coll Cardiol* 2006;113:2534-41
5. Cameli M, Ballo P, Righini FM, Caputo M, Lisi M, Mondillo S. Physiologic determinants of left ventricular systolic torsion assessed by speckle tracking echocardiography in healthy subjects. *Echocardiography* 2011;28:641–8.
6. Tan YT, Wenzelburger F, Lee E, Heatlie G, Leyva F, Patel K, et al: The pathophysiology of heart failure with normal ejection fraction: exercise echocardiography reveals complex abnormalities of both systolic and diastolic ventricular function involving torsion, untwist, and longitudinal motion. *J Am Coll Cardiol* 2009;54:36–46.
7. Saygısunar U, Kılıç H, Aytürk M, et al: Volume depletion provided by blood donation alters twist mechanics of the heart. *Scandinavian Cardiovascular Journal* 2016;50:23-27.
8. Russel IK, Gotte MJ, Bronzwaer JG, Knaapen P, Paulus WJ, van Rossum AC. Left ventricular torsion: an expanding role in the analysis of myocardial dysfunction. *JACC Cardiovasc Imaging* 2009;2:648–55.
9. Grabskaya E, Spira C, Hoffmann R, Altiok E, Ocklenburg C, Hoffmann R, et al. Myocardial rotation but not circumferential strain is transducer angle dependent: a speckle tracking echocardiography study. *Echocardiography* 2010;27:809-14.

10. Sengupta PP, Tajik AJ, Chandrasekaran K, Khandheria BK. Twist mechanics of the left ventricle: principles and application. *JACC Cardiovasc Imaging* 2008;1:366-76.
11. Sohn DW, Chai IH, Lee DJ, Kim HC, Kim HS, Oh BH, et al. Assessment of mitral annulus velocity by Doppler tissue imaging in the evaluation of left ventricular diastolic function. *J Am Coll Cardiol* 1997;30:474-80
12. Kim HK, Sohn DW, Lee SE, Choi SY, Park JS, Kim YJ, et al. Assessment of left ventricular rotation and torsion with two-dimensional speckle tracking echocardiography. *J Am Soc Echocardiogr* 2007;20:45-53.
13. Nakai H, Takeuchi M, Nishikage T, Kokumai M, Otani S, Lang RM. Effect of aging on twist-displacement loop by 2-dimensional speckle tracking imaging. *J Am Soc Echocardiography* 2006;19:880-5.
14. Helle-Valle T, Crosby J, Edvardsen T, Lyseggen E, Amundsen BH, Smith HJ, et al. New noninvasive method for assessment of left ventricular rotation: speckle tracking echocardiography. *Circulation* 2005; 112: 3149-56.
15. Notomi Y, Lysyansky P, Setser RM, Shiota T, Popovic ZB, Martin- Miklovic MG, et al. Measurement of ventricular torsion by two-dimensional ultrasound speckle tracking imaging. *J Am Coll Cardiol* 2005; 45: 2034-41.
16. Nagel E, Stuber M, Burkhard B, Fischer SE, Scheidegger MB, Boesiger P, et al. Cardiac rotation and relaxation in patients with aortic valve stenosis. *Eur Heart J* 2000; 21:582-9.
17. Tadic M, Cuspidi C, Majstorovic A, Kocijancic V, Celic V. The relationship between left ventricular deformation and different geometric patterns according to the updated classification: findings from the hypertensive population. *J Hypertens*. 2015;33:1954-61
18. Sandstede JJ, Johnson T, Harre K, Beer M, Hofmann S, Pabst T, et al. Cardiac systolic rotation and contraction before and after valve replacement for aortic stenosis: a myocardial tagging study using MR imaging. *AJR Am J Roentgenol* 2002;178:953-8.

19. Takeuchi M, Nakai H, Kokumai M, Nishikage T, Otani S, Lang RM. Age-related changes in left ventricular twist assessed by twodimensional speckle-tracking imaging. *J Am Soc Echocardiogr* 2006; 19:1077–84.
20. Sun JP, Lam YY, Wu CQ, Yang XS, Guo R, Kwong JS, et al. Effect of age and gender on left ventricular rotation and twist in a large group of normal adults: a multicenter study. *Int J Cardiol.* 2013; 167:2215-21
21. Zhang Y, Zhou QC, Pu D, Zou L, Tan Y. Differences in left ventricular twist related to age: speckle tracking echocardiographic data for healthy volunteers from neonate to age 70 years. *Echocardiography* 2010;27:1205–10.
22. Lee HJ, Kim BS, Kim JH, Jang HS, Bae BS, Kang HJ, et al. Age-related changes in left ventricular torsion as assessed by 2-dimensional ultrasound speckle tracking imaging. *Korean Circ J* 2008; 38:529–35.
23. Maufrais C, Schuster I, Doucende G, Vitiello D, Rupp T, Dautat M, et al. Endurance Training Minimizes Age-Related Changes of Left Ventricular Twist-Untwist Mechanics. *J Am Soc Echocardiogr* 2014;27:1208-15
24. Pagourelas ED, Sotiriou P, Papadopoulos CE, Cholongitas E, Giouleme O, Vassilikos V. Left Ventricular Myocardial Mechanics in Cirrhosis: A Speckle Tracking Echocardiographic Study. *Echocardiography.* 2015 Jul 14. doi: 10.1111/echo.13010.
25. Enache R, Popescu BA, Piazza R, Muraru D, Călin A, Beladan CC, et al. Left ventricular shape and mass impact torsional dynamics in asymptomatic patients with chronic aortic regurgitation and normal left ventricular ejection fraction. *Int J Cardiovasc Imaging.* 2015;31:1315-26.

Table 1: Baseline clinical characteristic values and echocardiographic findings of all subjects

	Young	Older	P value
Age (years)	29±6	64±12	0.001
Patients number (n/M:F)	39/17:22	36/14:22	0.596
Hypertension n(%)	1(2.6%)	17(48.6%)	0.000
diabetes mellitus (%)	1(2.6%)	6(17.1%)	0.03
BMI (kg/m ²)	24.5±4.8	29.6±5.2	0.001
LVEF (%)	69.2±7	65.9±7	0.045
E	76.8±11	52.4±16	0.000
A	57.7±10.6	70.6±13.7	0.000
E'	13.1±3.6	7±1.9	0.000
E/A	1.3	0.75	0.000
E/E' ratio	6.1±1.3	7.7±2.2	0.000
Apical rotation ,°	3.3±2.4°	4.4±2.9°	0.075
Basal rotation ,°	-5.2±3.7°	-4.1±2.7°	0.185
Twist ,°	8.5±4.1°	8.7±3.3°	0.890

A, late mitral inflow velocity; BMI, body mass index; E, early mitral inflow velocity; E', early diastolic mitral annular velocity; LVEF, left ventricular ejection fraction;

Table 2: Change in rotation and twist with diastolic dysfunction in old group

	Normal (7)	Grade 1 LVDD (23)	Grade 2 LVDD(6)
apical rotation, °,	4.1±2.5	5.2±3.2	2.4±1.2
basal rotation, °,	-4.9±-3.6	-4.4 ±-2.7	-4.5±-2.9
twist	6.9±2.4	9.7±3.2	6.7±3.3

LVDD, left ventricular diastolic dysfunction

Figure 1: Apical and basal rotation curves. The LV rotates counter-clockwise at the apical level and clockwise at the base. Counter-clockwise LV rotation as viewed from the apex is expressed as a positive value and clockwise LV rotation is expressed as a negative value.

