



# Comparison of Left Atrial Function in Long-Term Anabolic/Androgenic Steroid Users versus Nonuser Bodybuilders by Using Two-Dimensional Speckle-Tracking Echocardiography

Elnur Alizade

University of Health Sciences, Kartal Koşuyolu High Specialization Health Application and Research Center, Clinic of Cardiology, İstanbul, Turkey

## ABSTRACT

**Introduction:** Long-term illicit use of supraphysiological doses of anabolic/androgenic steroids (AAS) may cause pathological left ventricular hypertrophy (LVH), diastolic dysfunction, left atrial (LA) hypertrophy, increased myocardial stiffness, and myocardial fibrosis. Therefore, distinguishing AAS-using athlete's hearts from the nonpathological "athlete's heart" is critically important. The aim of this study was to evaluate LA myocardial function using 2D-STE method in both AAS-using and drug-free bodybuilders, and assess its potential role in the differential diagnosis between these two entities.

**Patients and Methods:** We selected a population of 33 male, competitive bodybuilders, including 15 actively using AAS for > 2 years (users) and 18 who had never used AAS (nonusers).

**Results:** AAS users had a significantly lower global LA strain reservoir (GLAS-R), global LA strain during early diastole (GLAS-E;  $38.2 \pm 8.4$  vs.  $48.6 \pm 11.9$ ,  $p < 0.01$ ;  $24.4 \pm 8.6$  vs.  $37.1 \pm 12.8$ ,  $p < 0.01$ ; respectively), global LA strain rate reservoir (GLASR-R), global LA strain rate during early diastole (GLASR-E;  $1.8 \pm 0.3$  vs.  $2.2 \pm 0.4$ ,  $p < 0.01$ ;  $-1.4 \pm 0.2$  vs.  $-1.8 \pm 0.3$ ,  $p < 0.01$ ; respectively) than nonusers. The univariate correlation analysis demonstrated that GLAS-R, GLAS-E, GLASR-R, and GLASR-E had a good inverse correlation with E/Em (r:  $-0.34$ ,  $p = 0.04$ ; r:  $-0.35$ ,  $p = 0.04$ ; r:  $-0.35$ ,  $p = 0.04$ , and r:  $-0.35$ ,  $p = 0.04$ , respectively).

**Conclusion:** The present study confirms that LA strain and strain rate are impaired in AAS users compared to nonusers and provide valuable additional information to that obtained by conventional echocardiography in the differential diagnosis between pathological and physiological LVH.

**Key Words:** Anabolic/androgenic steroid; bodybuilder athletes; left atrial function; speckle tracking; left atrial strain

## Uzun Dönem Anabolik-Androjenik Steroid Kullanan ve Kullanmayan Vücut Geliştiricilerde Sol Atrial Fonksiyonların İki Boyutlu Benekli İşaretleme Ekokardiyografik Yöntemi ile Karşılaştırılması

### ÖZET

**Giriş:** Uzun dönem ve yüksek dozda anabolik-androjenik steroid (AAS) kullanımı sol ventrikül patolojik hipertrofisine (SVH), diyastolik disfonksiyona, sol atrium (SA) hipertrofisine, miyokart sertlik ve fibrozisinde artmaya neden olur. Bundan dolayı AAS kullanan sporcuların patolojik olmayan "sporcu kalbi"nden ayrımı kritik öneme sahiptir. Bu çalışmanın amacı SA miyokart fonksiyonlarının iki boyutlu benekli ekokardiyografik yöntem ile AAS kullanan ve kullanmayan vücut geliştiricilerde değerlendirilmesi ve bu yöntemin bu iki durumun ayrımında kullanılmasıdır.

**Hastalar ve Yöntem:** Çalışmamıza 15'i aktif olarak AAS kullanan (> 2 yıl) ve 18'i hiç AAS kullanmayan hepsi erkek olan toplam 33 yarışmacı vücut geliştirici alındı.

**Bulgular:** AAS kullanan atletlerin global SA strain reservoir (GLAS-R), global SA erken diyastol strain (GLAS-E) ( $38.2 \pm 8.4$  vs.  $48.6 \pm 11.9$ ,  $p < 0.01$ ;  $24.4 \pm 8.6$  vs.  $37.1 \pm 12.8$ ,  $p < 0.01$ ; sırasıyla), global SA strain rate reservoir (GLASR-R), global SA erken diyastol strain rate (GLASR-E) ( $1.8 \pm 0.3$  vs.  $2.2 \pm 0.4$ ,  $p < 0.01$ ;  $-1.4 \pm 0.2$  vs.  $-1.8 \pm 0.3$ ,  $p < 0.01$ ; sırasıyla) değerlerinin kullanmayan sporculara göre önemli ölçüde azaldığı görüldü. Univaryant korelasyon analizi GLAS-R, GLAS-E, GLASR-R ve GLASR-E değerlerinin E/Em (r:  $-0.34$ ,  $p = 0.04$ ; r:  $-0.35$ ,  $p = 0.04$ ; r:  $-0.35$ ,  $p = 0.04$  and r:  $-0.35$ ,  $p = 0.04$ , sırasıyla) ile ters korelasyon olduğunu gösterdi.

**Sonuç:** Bu çalışma SA strain ve strain rate değerlerinin AAS kullanan sporcularda kullanmayan sporculara göre azaldığı, ayrıca patolojik ve fizyolojik SVH ayırıcı tanısında konvansiyonel ekokardiyografi ile elde edilen bulgulara ek değerli bilgiler sağladığı doğrulanmıştır.

**Anahtar Kelimeler:** Anabolik-androjenik steroid; vücut geliştirici sporcular; sol atrial fonksiyon; benekli işaretleme; sol atrial strain

## Correspondence

Elnur Alizade

E-mail: elnur17@yahoo.com

Submitted: 06.11.2016

Accepted: 29.11.2016

© Copyright 2017 by Koşuyolu Heart Journal.  
Available on-line at  
www.kosuyoluheartjournal.com

## INTRODUCTION

Self-administration of high doses of anabolic/androgenic steroids (AAS) is a widespread practice among athletes to increase lean body mass and muscular strength. Long-term illicit use of supraphysiological doses of AAS may cause several adverse cardiovascular effects<sup>(1-4)</sup>. There are several case reports of sudden death in athletes, which indicates an association between chronic AAS abuse and increased risk of arrhythmias and sudden cardiac death<sup>(5,6)</sup>. It has been reported that cardiovascular morbidity and mortality have significantly increased in long-term AAS using bodybuilders than nonusers<sup>(5)</sup>. Furthermore, recent studies have found pathological left ventricular hypertrophy (LVH), diastolic dysfunction [impaired relaxation and decreased compliance of left ventricle (LV)], increased LV mass, left atrial (LA) hypertrophy, subclinical systolic impairment, increased myocardial stiffness and myocardial fibrosis in long-term AAS users<sup>(4,7-9)</sup>. These conditions are independent risk factors for cardiovascular morbidity and mortality. Therefore, distinguishing AAS users' heart from the nonpathological "athlete's heart" is critically important. Echocardiography plays a key role in differential diagnosis, but significant overlap exists between both conditions and its differentiation remains challenging.

LA volume and function are useful barometers of LV diastolic function and predictors of cardiovascular outcomes.<sup>(10)</sup> Changes in LA size or phasic function may indicate the presence and the severity of heart disease<sup>(11,12)</sup>. Pathological LVH is associated with LA dilatation and dysfunction. Echocardiographic assessment of LA myocardial function might contribute to differentiate between pathological and physiological LVH<sup>(13)</sup>. Recently, two-dimensional S and SR echocardiographic imaging based on 2D-STE have been proposed as a method for assessing LA function. 2D-STE is a novel non-Doppler-based method for the angle-independent and objective quantification of myocardial deformation from bidimensional datasets in contrast to Doppler-derived indices. In addition, speckle tracking has the advantage of being angle independent and being less affected by reverberations, side lobes and dropout artifacts<sup>(14)</sup>. This analysis may allow a more direct assessment of LA endocardial contractility, and passive deformation has been recently proposed. The feasibility and reproducibility of 2D-STE for the study of LA mechanics have been recently validated. Abnormalities in LA S and SR have been shown by 2D-STE in some pathophysiological conditions, including systolic and diastolic heart failure, atrial fibrillation, stroke, heart valve disease and hypertrophic cardiomyopathy<sup>(14,15)</sup>.

The LA function contributes to LV filling by means of its three components: a reservoir component, which receives blood from the pulmonary veins during ventricular systole; a

passive conduit component during early diastole and diastasis; and a pump component, with active contraction during late diastole. LV dysfunction, which arises in pathologies that affect LV structure and function, affects LA functions; in addition, LA functions can be affected in relation to the negative impact of these pathologies on the structure of the LA<sup>(15)</sup>.

Although LV systolic and diastolic functions have been examined in various echocardiographic researches, no research has been conducted to examine LA functions between AAS users and nonusers using the 2D-STE method. The aim of this study was to evaluate LA myocardial function using 2D-STE method in both AAS-using bodybuilders and drug-free bodybuilders, and assess its potential role in the differential diagnosis between these two entities.

## PATIENTS and METHODS

### Study Population

We selected a population of 33 competitive bodybuilders, including 15 actively using AAS for  $\geq 2$  years (users) and 18 who had never used AAS (nonusers), all men. The protocol of this cross-sectional study was approved by the institutional review board of Kartal Kosuyolu Heart and Training Hospital and performed in accordance with the guidelines proposed in the Helsinki Declaration. Written informed consent was obtained from all the participants. Patients with coronary artery disease, chronic renal failure, chronic liver disorders, chronic lung disease, moderate or severe valvular heart disease, diabetes mellitus, congenital heart disease, LV systolic dysfunction on echocardiography (EF < 50%), recent acute coronary syndrome, anemia, obstructive sleep apnea, secondary hypertension, hematological disorders, known malignancy, thyroid dysfunction, hypercholesterolemia, electrolyte imbalance and bundle-branch block, atrioventricular conduction abnormalities on ECG were excluded from the study. All the patients were in sinus rhythm and none of them was taking medications such as antiarrhythmics, tricyclic antidepressants, antihistamine agents, and antipsychotics.

**Training protocols:** All participants had trained intensively for > 10-15 h/week for > 5 years. AAS users and nonusers had started bodybuilding at approximately the same age ( $21.47 \pm 3.24$  vs.  $22.34 \pm 3.68$  years, respectively,  $p =$  nonsignificant) and training protocols (anaerobic isometric static exercises and aerobic exercises) were not different between the groups. Maximum self-reported one-repetition squat results were significantly greater among AAS users ( $142.6 \pm 19.0$  vs.  $120.6 \pm 21.6$  kg,  $p < 0.001$ ; Table 1).

**AAS abuse:** A self-reported clinical history of each participant including type and timing of AAS use and other performance-enhancing drugs was carefully noted. The orally

**Table 1. Training programs of the AAS users and nonusers**

	AAS nonusers (n= 18)	AAS users (n= 15)	p
Sessions per week	3.92 ± 0.86	3.67 ± 0.84	NS
Years	8.64 ± 2.11	9.03 ± 1.94	NS
Starting age	22.34 ± 3.68	21.61 ± 3.04	NS
Anaerobic exercise (h/week)	4.73 ± 2.02	4.94 ± 1.82	NS
Aerobic exercise (h/week)	3.11 ± 3.03	1.94 ± 1.82	NS
Maximal weight (kg)	120.67 ± 21.61	142.67 ± 19.09	< 0.05

AAS: Anabolic/androgenic steroids, NS: Nonsignificant.

self-administered drugs were oxymetholone and stanozolol, and the injectable steroids were nandrolone, stanozolol, and testosterone propionate. The mean duration of AAS use was  $5.73 \pm 3$  years. The mean weekly dosage of AAS was  $1085.5 \pm 354$  mg.

**Physical examination and laboratory tests:** All subjects were examined on an empty stomach. Height, weight, body mass index, body surface area (BSA), body fat mass, heart rate, and blood pressure were measured. Venous blood samples were drawn from each subject, always in the afternoon between 1 and 2 PM, to evaluate serum hormone levels (testosterone, luteinizing hormone, follicle-stimulating hormone, insulin, T3, and T4), hematology (hematocrit, hemoglobin), and blood lipids (total cholesterol, high-density lipoprotein). The subjects' bodyweight and height were measured and the body mass index (BMI) was calculated as body weight divided by squared height ( $\text{kg}/\text{m}^2$ ).

#### Echocardiographic Measurements

Echocardiography was performed in left lateral decubitus position with an ultrasound machine GE Vingmed Vivid 7 system (Vivid system 7, GE Vingmed Ultrasound AS, Horten, Norway) and 3S-RS (3.5 MHz) probe. Examinations were performed by a cardiologist who was blinded to the clinical details of each subject. Single-lead ECG was recorded continuously during the echocardiographic examination. Two-dimensional, M-mode, and tissue Doppler images were acquired from the parasternal long-axis and short-axis and apical four-chamber views at end-expiratory apnea, and were transferred to a customized dedicated software package (EchoPAC, General Electric Vingmed Ultrasound) for offline analysis of stored data. All measurements were averaged from three cardiac cycles. 2D echocardiographic measurements were performed according to standards outlined by the American Society of Echocardiography<sup>(16)</sup>.

#### LV Assessment

LV dimensions and wall thickness were obtained from the parasternal long axis with an M-mode cursor positioned just beyond the mitral leaflet tips, perpendicular to the long

axis of the ventricle. LV end-diastolic diameter (LVEDD) and end-systolic (LVESD) diameter, thickness of the interventricular septum (IVS), and the LV posterior wall (PW) were measured. LV ejection fraction was calculated according to the Simpson method<sup>(16)</sup>. For determination of LV mass (LVM), the Devereux formula was used:  $\text{LVM (g)} = 1.04 [(\text{LVID} + \text{PWT} + \text{IVST})^3 - \text{LVID}^3] - 14$  (LVID indicates LV internal dimension; PWT, PW thickness; IVST, IVS. thickness). LV mass index was calculated by dividing LVM by BSA. LV hypertrophy was defined as an LV mass index  $> 115 \text{ g}/\text{m}^2$  in men, as recommended by the American Society of Echocardiography and the European Association of Echocardiography<sup>(16)</sup>. Mitral inflow velocities were evaluated by pulsed-wave Doppler echocardiography with the sample volume placed at the tip of the mitral leaflets from the apical four-chamber view. Diastolic peak early (E) and peak late (A) transmittal flow velocity, peak E to peak A velocities (E/A), deceleration time of peak E velocity (EDT) and isovolumetric relaxation time (IVRT) were measured<sup>(17)</sup>.

The tissue Doppler imaging (TDI) was performed in the apical four-chamber view using a 5-mm pulsed-wave Doppler sample volume with as minimum optimal gain as possible to obtain the best signal-to-noise ratio. Care was taken to align the echo image so that the annular motion was parallel to the TDI cursor. Spectral pulsed-wave Doppler signal filters were adjusted until a Nyquist limit of 15-20 cm/s was reached. The monitor sweep speed was set at 50-100 mm/s to optimize the spectral display of myocardial velocities. In apical four-chamber view, the pulsed-wave Doppler sample volume was subsequently placed at the level of LV lateral mitral annulus, septal mitral annulus, and right ventricular (RV) tricuspid annulus. The myocardial peak systolic (Sm), and early diastolic (Em) velocity, and late diastolic (Am) velocity were obtained from the septum, the lateral wall of LV and RV annulus. The Sm global, Em global, and Am global velocities were derived by averaging the velocities from the 2 mitral annular sites. Global Em/Am ratio and E/Em ratio were calculated<sup>(18)</sup>.


### LA Assessment

All volumes were calculated from the apical four-chamber and two-chamber views using the Simpson biplane method of discs<sup>(19)</sup>. LA length was defined as the longest line that could be drawn between the posterior LA wall and the mid-portion of the mitral valve, and was similar in the four-chamber and two-chamber views, which are perpendicular to each other. Maximum LA volume was measured just before mitral valve opening. Minimum LA volume was measured at mitral valve closure. All volumes were indexed to BSA. LA volume at onset of atrial systole was considered the volume corresponding to the onset of the P wave in the simultaneously recorded ECG. All LA volume values were corrected for BSA. LA systolic (active emptying) function was assessed using (1) LA active emptying volume = LA volume at onset of atrial systole LA minimal volume and (2) LA active emptying fraction = LA active emptying volume/LA volume at onset of atrial systole<sup>(20,21)</sup>.

To analyze 2D speckle-tracking imaging, we obtained 2D gray scale harmonic images from the apical four-chamber and two-chamber views focused on the LA. All images were obtained at a frame rate of 40-90 frames/s without dual focus. Three consecutive cardiac cycles were saved in digital format for offline analysis using dedicated software (EchoPAC version 8.0.0; GE Vingmed). The software is based on real-time tracking of natural acoustic markers, which allows the derivation of 2D strain and strain rate by comparing the

relative displacement of speckles throughout the cardiac cycle. To determine the LA longitudinal strain (S) and strain rate (SR), the endocardial border of the LA was traced manually and tracked by the software. The software divided each LA wall arbitrarily into three segments: annular, mid, and basal segments. In the case of unsatisfactory tracking, the operator could repeat the imaging or change software parameters such as the region of interest width and the smoothing functions. When more than two inadequately traced segments were found among six segments in each apical view even after the repeated tracing, the image was excluded from the analysis. Inadequately traced segments were automatically excluded from analysis. Finally, the software calculated average SR for six segments for each apical view and the LA S-SR values for each view were the averages of the values obtained for the LA segments of each view. The final Global LA-S and -SR values were the averages of the values obtained for each apical view<sup>(22,23)</sup>.

LA strain during systole (LAS-S) was obtained at the time of aortic valve closure, strain during late diastole (LAS-A) was obtained at the onset of the P wave on electrocardiography, peak systolic strain rate (LASR-S), peak early diastolic strain rate (LASR-E) and peak late diastolic strain rate (LASR-A) were obtained for the entire traced contour of the LA. LA strain during early diastole (LAS-E) was defined as (LAS-S)–(LAS-A). LAS-S and LASR-S LA reservoir, LAS-E and LASR-E LA conduit, LAS-A and LASR-A values are related to pump functions of the LA<sup>(23)</sup> (Figure 1).



**Figure 1.** (A) Left atrial global strain reservoir (GLAS-R), global strain early diastole (GLAS-E), and global strain late diastole (GLAS-A), (B) left atrial global strain rate reservoir (GLASR-R), global strain rate early diastole (GLASR-E), and global strain rate late diastole (GLASR-A).



### Statistical Analysis

The SPSS 15.0 statistical program (SPSS Inc., Chicago, Ill.) was used for the statistical analysis. All values are given as means  $\pm$  standard deviations. Mean values of continuous variables were compared between the groups using the Student t-test or Mann-Whitney U-test, according to whether normally distributed or not, as tested by the Kolmogorov-Smirnov test. The chi-square test was used to assess differences between categorical variables. Spearman's correlation coefficients were used to assess the strength of relationship between continuous variables. A p value of  $< 0.05$  was considered significant.

## RESULTS

### Clinical Characteristics of the Study Population

The characteristics of the subjects are listed in Table 2. No differences between the groups emerged in age, height, weight, BSA, blood pressure, or heart rate. However, AAS users had higher body mass indices than AAS nonusers.

### Echocardiographic Analysis

Table 3 shows the details of the echocardiographic analysis. LV mass index, interventricular septal thickness, LV posterior wall thickness, and relative diastolic wall thickness were significantly greater in AAS users than in nonusers ( $p < 0.01$ ). No significant differences were found in LV end-systolic dimension, end-diastolic dimension, and ejection fraction among the groups.

Transmitral Doppler echocardiography data of LV diastolic function are listed in Table 3. No significant differences were found in peak E and peak A between AAS users and nonusers. However, drug-using bodybuilders exhibited longer isovolumetric relaxation times and lower ratio of E/A than their drug-free counterparts.

When comparing the diastolic functions obtained by measuring the TDI velocities, lateral, and septal Em were significantly lower in AAS users than in nonusers ( $11.6 \pm 1.2$

vs.  $16.2 \pm 1.5$ ,  $p < 0.01$ ;  $10.1 \pm 1.3$  vs.  $12.1 \pm 1.5$ ,  $p < 0.01$ ; respectively), whereas lateral Am and septal Am were not a significant difference in AAS users than in nonusers ( $9.4 \pm 1.3$  vs.  $9.9 \pm 1.2$ ,  $p > 0.05$ ;  $9.5 \pm 0.7$  vs.  $9.4 \pm 1.2$ ,  $p > 0.05$ ; respectively). Global E/Em and Em/Am were significantly different in AAS users than in nonusers ( $7.3 \pm 1.5$  vs.  $5.8 \pm 0.9$ ,  $p < 0.01$ ;  $1.6 \pm 0.1$  vs.  $1.5 \pm 0.2$ ,  $p < 0.01$ ; respectively). In addition, Sm was significantly lower in AAS users than in nonusers ( $6.23 \pm 0.63$  vs.  $7.04 \pm 1.16$ ,  $p < 0.01$ ).

LA active emptying volume and active emptying fraction were increased in AAS users than in nonusers ( $4.6 \pm 1.0$  vs.  $3.07 \pm 1.1$ ,  $p < 0.001$ ;  $33.1 \pm 8.1$  vs.  $25.8 \pm 9.1$ ,  $p < 0.001$ ; respectively), but LA volume index was not different between these groups. With respect to the LA strain and strain rate, AAS using athletes had significantly lower Global LAS-R, Global LAS-E ( $38.2 \pm 8.4$  vs.  $48.6 \pm 11.9$ ,  $p < 0.01$ ;  $24.4 \pm 8.6$  vs.  $37.1 \pm 12.8$ ,  $p < 0.01$ ; respectively), Global LASR-R, Global LASR-E ( $1.8 \pm 0.3$  vs.  $2.2 \pm 0.4$ ,  $p < 0.01$ ;  $-1.4 \pm 0.2$  vs.  $-1.8 \pm 0.3$ ,  $p < 0.01$ ; respectively) than nonusers. Global LAS-A and Global LASR-A was also increased in AAS users (Table 4).

The univariate correlation analysis demonstrated that the Global LAS-R, Global LAS-E, Global LASR-R, and Global LASR-E had a good inverse correlation with E/Em ( $r: -0.34$ ,  $p = 0.04$ ;  $r: -0.35$ ,  $p = 0.04$ ;  $r: -0.35$ ,  $p = 0.04$ , and  $r: -0.35$ ,  $p = 0.04$ , respectively). There were also negative correlation between Global LAS-R, Global LAS-E, Global LASR-R, Global LASR-E, and LV mass index ( $r: -0.38$ ,  $p = 0.02$ ;  $r: -0.39$ ,  $p = 0.02$ ;  $r: -0.37$ ,  $p = 0.03$ , and  $r: -0.40$ ,  $p = 0.02$ , respectively). LA strain and strain rate parameters all had a positive correlation with both Em and Sm (Table 5).

Intraobserver and interobserver variability coefficients were calculated using images independently recorded at two different occasions by the same investigator or by two different observers. When the reproducibility was separately considered in the two apical views, interobserver variability coefficients were 5.4% and 5.9% for four-chamber and two-chamber average Global LAS-R, 6% and 6.1% for Global LASR-R,

**Table 2. Clinical characteristics of AAS users and nonusers**

	AAS nonusers (n= 18)	AAS users (n= 15)	p
Age (year)	33.8 $\pm$ 4.1	32.5 $\pm$ 6.6	NS
Height (cm)	180.4 $\pm$ 6.9	179.9 $\pm$ 7.3	NS
Weight (kg)	87.4 $\pm$ 10.3	90.8 $\pm$ 6.3	NS
BMI (kg/m <sup>2</sup> )	26.3 $\pm$ 3.2	29.1 $\pm$ 4.4	< 0.05
BSA (m <sup>2</sup> )	2.08 $\pm$ 0.14	2.1 $\pm$ 0.14	NS
Blood pressure (mmHg)	120 $\pm$ 13.37    80.37 $\pm$ 6.49	118.51 $\pm$ 9.88    78.51 $\pm$ 6.9	NS
Heart rate (beats/min)	68.74 $\pm$ 10.45	72.22 $\pm$ 13.40	NS

AAS: Anabolic/androgenic steroids, NS: Nonsignificant, BMI: Body mass index, BSA: Body surface area.

**Table 3. Comparison of the echocardiographic parameters of both AAS users and nonusers**

	AAS nonusers (n= 18)	AAS users (n= 15)	p
<b>2-D Echocardiographic parameters</b>			
LV end-systolic diameter (mm)	31.9 ± 4.4	33.2 ± 3.2	NS
LV end-diastolic diameter (mm)	49.7 ± 1.9	51.2 ± 3.1	NS
Septal wall thickness (mm)	11.5 ± 1.2	12.4 ± 1.3	< 0.01
Posterior wall thickness (mm)	9.8 ± 0.9	11.3 ± 0.7	< 0.01
RWT	0.39 ± 0.03	0.44 ± 0.02	< 0.01
LV mass index (g/m <sup>2</sup> )	90.9 ± 10.8	113.6 ± 13.6	< 0.01
LV ejection fraction (%)	61.37 ± 1.6	60.87 ± 2.3	NS
<b>Doppler parameters</b>			
Peak E velocity (cm/s)	79.8 ± 9.4	77.6 ± 11.6	NS
Peak A velocity (cm/s)	55.7 ± 8.9	50.7 ± 6.8	NS
E/A ratio	1.47 ± 0.3	1.54 ± 0.2	NS
IVRT (ms)	80.7 ± 5.8	83.58 ± 11.7	< 0.01
S <sub>m</sub> (cm/s)	7.04 ± 1.16	6.23 ± 0.63	< 0.01
E <sub>m</sub> septal (cm/s)	12.1 ± 1.5	10.1 ± 1.3	< 0.01
A <sub>m</sub> septal (cm/s)	9.4 ± 1.2	9.5 ± 0.7	NS
E/E <sub>m</sub> septal (cm/s)	6.7 ± 1.2	7.8 ± 1.7	< 0.01
E <sub>m</sub> /A <sub>m</sub> septal (cm/s)	1.29 ± 0.2	1.06 ± 0.2	< 0.01
E <sub>m</sub> lateral (cm/s)	16.2 ± 1.5	11.6 ± 1.2	< 0.01
A <sub>m</sub> lateral (cm/s)	9.9 ± 1.2	9.4 ± 1.3	NS
E/E <sub>m</sub> lateral (cm/s)	4.9 ± 0.8	6.8 ± 1.3	< 0.01
E <sub>m</sub> /A <sub>m</sub> lateral (cm/s)	1.6 ± 0.3	1.2 ± 0.2	< 0.01
E/E <sub>m</sub> global (cm/s)	5.8 ± 0.9	7.3 ± 1.5	< 0.01
E <sub>m</sub> /A <sub>m</sub> global (cm/s)	1.5 ± 0.2	1.6 ± 0.1	< 0.01

AAS: Anabolic/androgenic steroids, NS: Nonsignificant, LV: Left ventricle, RWT: Relative wall thickness, IVRT: Isovolumetric relaxation time, S<sub>m</sub>: Peak systolic myocardial velocity, E<sub>m</sub>: Early diastolic myocardial velocity, A<sub>m</sub>: Late diastolic myocardial velocity.

**Table 4. Left atrial standard echocardiography and two-dimensional strain baseline measurements in the overall study population**

	AAS nonusers (n= 18)	AAS users (n= 15)	p
LA dimension (mm)	33.1 ± 0.3	34.2 ± 0.2	NS
LA volume index (mL/m <sup>2</sup> )	26.2 ± 2.3	27.6 ± 2.4	NS
LA active emptying volume (cm <sup>2</sup> /m <sup>2</sup> )	3.07 ± 1.1	4.6 ± 1.0	< 0.01
LA active emptying fraction (%)	25.8 ± 9.1	33.1 ± 8.1	0.02
GLAS-R (%)	48.6 ± 11.9	38.2 ± 8.4	< 0.01
GLAS-E (%)	37.1 ± 12.8	24.4 ± 8.6	< 0.01
GLAS-A (%)	11.5 ± 2.4	13.8 ± 2.2	< 0.01
GLASR-R (1/s)	2.2 ± 0.4	1.8 ± 0.3	< 0.01
GLASR-E (1/s)	-1.8 ± 0.3	-1.4 ± 0.2	< 0.01
GLASR-A (1/s)	-1.4 ± 0.2	-1.6 ± 0.2	0.02

AAS: Anabolic/androgenic steroids, NS: Nonsignificant, LA: Left atrium, GLAS-R: Global left atrial strain reservoir, GLAS-E: Global left atrial early diastolic strain, GLAS-A: Global left atrial late diastolic strain, GLASR-R: Global left atrial strain rate reservoir, GLASR-E: Global left atrial early diastolic strain rate, GLASR-A: Global left atrial late diastolic strain rate.

**Table 5. Spearman's correlation analysis (r and p values) between la strain/strain rate and echocardiographic parameters**

	E/E <sub>m</sub> global (cm/s)		LV mass index (g/m <sup>2</sup> )		Sm		E m (cm/s)	
	r (coefficient)	p	r (coefficient)	p	r (coefficient)	p	r (coefficient)	p
GLAS-R (%)	-0.34	<b>0.04</b>	-0.38	<b>0.02</b>	0.44	<b>0.01</b>	0.43	<b>0.01</b>
GLAS-E (%)	-0.35	<b>0.04</b>	-0.39	<b>0.02</b>	0.40	<b>0.02</b>	0.42	<b>0.01</b>
GLAS-A (%)	0.09	0.60	0.16	0.37	-0.13	0.46	-0.08	0.64
GLASR-R (%)	-0.35	<b>0.04</b>	-0.37	<b>0.03</b>	0.41	<b>0.01</b>	0.42	<b>0.01</b>
GLASR-E (%)	-0.35	<b>0.04</b>	-0.40	<b>0.02</b>	0.39	<b>0.02</b>	-0.41	<b>0.01</b>
GLASR-A (%)	-0.18	0.31	-0.27	0.11	0.21	0.22	0.16	0.37

GLAS-R: Global left atrial strain reservoir, GLAS-E: Global left atrial early diastolic strain, GLAS-A: Global left atrial late diastolic strain, GLASR-R: Global left atrial strain rate reservoir, GLASR-E: Global left atrial early diastolic strain rate, GLASR-A: Global left atrial late diastolic strain rate.

6.1% and 6% for Global LASR-E, 5.4% and 5.1% for Global LASR-A, respectively. Corresponding intraobserver variability coefficients were 4.7% and 5.1% for Global LAS-R, 5.4% and 5.6% for Global LAS-R, 5.4% and 5.2% for Global LASR-E, and 4.4% and 4.3% for Global LASR-A.

## DISCUSSION

The principal finding of this study is that assessment of LA strain and strain rate by 2D speckle tracking provides additional information to that obtained with standard echocardiography in the differential diagnosis between AAS using bodybuilders compared with non-AAS users. Furthermore, LA reservoir, conduit, and pump function are impaired in AAS users.

Hemodynamic overload due to long-term training usually involves both LV and RV, inducing changes in cardiac structure such as increases in internal cavity diameters, wall thickness, and mass, usually described as "athlete's heart"<sup>(24-26)</sup>. This is thought to be a physiological and benign response to exercise conditioning, without the adverse prognostic implications of HCM or hypertensive LV hypertrophy. The physiological hypertrophy that occurs in athletes is associated with increased ventricular mass, normal organization of cardiac structure, and no increase in collagen content. On the other side, the hypertrophy observed in AAS using bodybuilders is characterized by myofibrillar disarray and collagen accumulation, eventually leading to systolic and diastolic dysfunction<sup>(27,28)</sup>.

Echocardiography plays a key role in the early differential diagnosis between these two entities. Echocardiographic studies show that concentric LVH is a common morphological change of the heart in both long-term use of supraphysiological doses of AAS users and nonuser bodybuilder athletes. There are some studies related to the differences between pathological and physiological adaptive LVH in AAS using and nonusers<sup>(7,8)</sup>. In contrast to the physiological LVH caused by endurance training, LVH in pathological conditions like systemic hypertension

and hypertrophic cardiomyopathy is characterized by impaired diastolic function<sup>(29-31)</sup>. In addition, pathological LVH with impaired diastolic function, which is induced by long-term illicit use of supraphysiological doses of AAS has also been described<sup>(9,32,33)</sup>. In our study, we found that the E/Em ratio was significantly higher in AAS users than in nonusers. In addition, the Em/Am ratio was significantly lower in AAS users than in nonusers. In addition, we found that IVRT is prolonged in AAS users, indicating the impairment of diastolic function.

Although standard Doppler echocardiography has been widely used to distinguish athlete's heart from pathological LV hypertrophy, recent studies indicate that Doppler tissue imaging techniques, in particular PW-TDI, are also useful in assessing myocardial systolic and diastolic function and differentiating pathological ventricular hypertrophy from the physiological one<sup>(7,34,35)</sup>. Shan et al, comparing Doppler tissue imaging and histological findings in patients affected by coronary artery disease, demonstrated that Sm and Em are strongly dependent on the number of myocytes, myocardial b-adrenergic receptor density, and the amount of interstitial fibrosis<sup>(36)</sup>. D'Andrea et al. and other studies observed lower myocardial early diastolic peak velocities of the interventricular septum and the lateral LV wall in AAS users when than in nonusers<sup>(7,8)</sup>. Confirming previous findings, we observed low Sm velocity and low early diastolic peak velocities (Em) at the interventricular septum and the lateral LV wall, which indicate pathological LVH in AAS users.

LA function is an integral part of cardiac function that is often neglected. In recent years, LA strain and strain rate analysis by two-dimensional (2D) speckle tracking has emerged as a novel method to evaluate LA function. This analysis may allow a more direct assessment of LA endocardial contractility, and passive deformation has been recently proposed. The feasibility and reproducibility of 2D-STE for the study of LA mechanics have been recently validated. Abnormalities in LA-S and SR have been shown by STE in some pathophysiological

conditions, including systolic and diastolic heart failure, atrial fibrillation, stroke, and heart valve disease<sup>(14,37)</sup>. In addition, some authors have recently applied 2D strain echocardiography analysis to characterize LA myocardial function in patients with either physiological or pathological LVH<sup>(38)</sup>. It is known that hypertensive LVH shares many features with HCM and it has been suggested that evaluation of LA function may be relevant in the diagnosis of pathological LVH and in this sense LA-S and SR showed an acceptable reproducibility to assess LA function similar to other reports<sup>(13,39)</sup>. D'Andrea et al. found that LA myocardial deformation assessed by 2D speckle tracking is impaired in patients with LVH due to hypertension compared with elite athletes and control subjects<sup>(13)</sup>. In another study by Sun et al. demonstrated that all components of LA-SR in the hypertensive group were significantly lower than in athletes and normal controls<sup>(40)</sup>. Previous study showed that LA-S and SR were significantly decreased in patients with HCM than in highly trained athletes and healthy controls and provided valuable additional information to that obtained by conventional echocardiography in the differential diagnosis between pathological and physiological LVH<sup>(28)</sup>. Moreover, recent studies have found pathological LVH in long-term AAS users. Although conventional echocardiography has been widely used to distinguish pathological and physiological LVH between AAS users and nonusers, no research has been conducted to examine LA functions using the 2D-STE method between these groups. Thus, in the current study we evaluate LA myocardial function using 2D-STE method in both AAS users and nonusers to distinguish pathological and physiological LVH. In our study, in AAS users, Global LAS-A, Global LASR-A, and indices of LA booster pump function (LA active emptying fraction) were increased, whereas Global LA S/SR parameters, which are related with LA reservoir, conduit functions were significantly impaired compared with the nonusers. In addition, we found a correlation between Global LA S/SR reservoir and conduit functions with LV mass index, E/Em, Sm, and Em (Table 5). There was no correlation between Global LAS-A, Global LASR-A with LV mass index, E/Em, Sm, and Em (Table 5). Thus, we observed that Global LA-S and SR parameters give easy tool to differentiate pathological and physiological LVH between AAS users and nonusers. Global LA reservoir, conduit and pump functions in AAS users are impaired, even in the absence of LA enlargement.

There are several possible explanations for impaired Global LA-S and SR parameters in athletes with AAS users. Impaired LV diastolic function may be an early symptom of pathological hypertrophy and is helpful to differentiate between pathological and physiological LVH. The diastolic function may be restricted in AAS users, which has been described by our working group and others<sup>(7,8,41)</sup>. In our study, we described diastolic dysfunction in AAS using athletes. It is known that LA functions reflect the

severity of diastolic dysfunction. 2DSTE is a new modality that is capable of measuring phasic changes in LA strain<sup>(39,42)</sup>. It is known that LA reservoir and conduit function progressively declines at the advanced stages of diastolic dysfunction. This was associated with an initial augmentation of LA booster function in mild diastolic dysfunction to maintain total LA emptying volumes<sup>(43)</sup>. We found that Global LASR-A increased and Global LASR-E/R decreased with the development of LV diastolic dysfunction. The correlation of E/Em parameter with the Global LASR-E and Global LASR-R parameters suggests that LV diastolic dysfunction in AAS using athletes should have an impact on LA strain rate in early stage.

The other possible mechanism for impaired LA strain and strain rate parameters in AAS using athletes is LA myocardial fibrosis. Probably adverse effects of AAS on the cardiovascular system are also because of direct toxicity on myocardial structure with increased collagen deposition, fibrosis, and altered microcirculation with intimal hyperplasia of the intramural coronary arteries resulting in chronic ischemic damage<sup>(44)</sup>. As the cause of these alterations may directly affect the atrium and ventricle, causing inhomogeneous myocardial hypertrophy, focal myocyte damage with myofibrillar loss, and interstitial fibrosis heterogeneity in AAS users<sup>(44,45)</sup>. It is well known that LV interstitial fibrosis as a consequence of long-term AAS using in athletes, affects primarily the subendocardial systolic and diastolic function of LV<sup>(7,13,46)</sup>. In this regard, previous studies hypothesized that in long-term AAS using athletes, the same fibrotic processes that affect the subendocardial layer of LV could also alter the subendocardial fibers of the LA<sup>(44,45)</sup>. Several and recent studies suggest that the degree of elevated LV filling pressures may not fully explain LA failure and that LA myocardial fibrosis may play a role in the systolic and diastolic dysfunction of the LA. Systolic and diastolic myocardial functions of the LA are disrupted as a result of the fibrosis that occurs in the LA<sup>(47)</sup>. In pathologies that affect the LA structure and functions, the severity of the LA fibrosis evaluated with the magnetic resonance imaging as well as the correlation between the LA longitudinal strain and strain rate values proves the relationship between involved parameters and LA fibrosis<sup>(22)</sup>. The extent of atrial fibrosis detected with late gadolinium enhancement by magnetic resonance correlates with this reduction in atrial S and SR measured with speckle tracking<sup>(48)</sup>. In addition, several studies suggest that LA myocardial fibrosis play a role in the impairment of LA pump, conduit and reservoir functions in different cardiovascular diseases<sup>(48,49)</sup>. Suman et al. found that LA-S and SR (reservoir, conduit) were decreased in atrial fibrillation patients. In addition, they suggested that LA wall fibrosis by delayed-enhancement MRI is inversely related to LA-S and SR, and these are related to the AF burden<sup>(49)</sup>. Hence, we believe that the fibrosis that occurs in the LA due to long-term illicit use of supraphysiological doses of AAS might



be one of the reasons for the impairment of LA S/SR reservoir and conduit functions in AAS using bodybuilders.

Consequently, as described different cardiovascular diseases, we hypothesized that; LA myocardial fibrotic alterations, together with chronically elevated LV filling pressures (LV diastolic dysfunction), would lead to the decrease of LA reservoir and conduit function and to the increase of LA pump function (Frank-Starling law) in the long-term AAS users.

#### Study Limitations

The important limitation of our study is the small sample size, which makes necessary to assess the reproducibility of these results in larger scale studies. Our sample might not be representative of the overall population of long-term AAS users or weightlifters. The cross-sectional nature of this exploratory study does not permit to draw definitive conclusions about the long-term clinical implications of our findings.

We were dealing with young individuals. The impact of AAS in older individuals is unknown. The same idea can be used for gender. There is no guarantee that the effects of AAS on LA functions in women will be similar to those found in men. The information about the intake of steroids was self-reported, but it is difficult to assess this in an objective manner. In addition, training-related influences are also improbable as an explanation for the differences between the AAS users and nonusers in our study, as the training protocol was the same for all the athletes.

Cardiac MRI could not be performed. Therefore, LA remodeling and fibrosis could not be directly evaluated. The absence of studies in which LA deformation parameters obtained by 2D-STE were compared with sonomicrometry or tagged magnetic resonance imaging is another limitation. However, LV deformation parameters by 2D-STE are in good agreement with that obtained by sonomicrometry and by tagged magnetic resonance imaging<sup>(50)</sup>.

Finally, because a dedicated software for LA strain analysis has not been released yet, we used the current software for LV analysis to study the LA pattern strain. Future evolutions in this regard may be useful to improve tracking quality of LA myocardial deformation, and to provide a better instrument for the study of LA function.

#### CONCLUSION

The present study confirms that in AAS users, 2D-STE is effective and reliable noninvasive diagnostic tools for detecting early abnormalities of LA myocardial functions. In addition, this study shows that LA strain and strain rate are impaired in AAS users compared to nonusers and provide valuable additional information to that obtained by conventional echocardiography in the differential diagnosis between pathological and physiological LVH

#### CONFLICT of INTEREST

The authors reported no conflict of interest related to this article.

#### AUTHORSHIP CONTRIBUTIONS

*Concept/Design:* EA

*Analysis/Interpretation:* EA

*Data Acquisition:* EA

*Writing:* EA

*Critical Revision:* EA

*Final Approval:* EA

#### REFERENCES

1. Kanayama G, Hudson JI, Pope HG Jr. Long-term psychiatric and medical consequences of anabolic-androgenic steroid abuse: a looming public health concern? *Drug Alcohol Depend* 2008;98:1-12.
2. Parssinen M, Seppala T. Steroid use and long-term health risks in former athletes. *Sports Med* 2002;32:83-94.
3. Thiblin I, Petersson A. Pharmacoepidemiology of anabolic androgenic steroids: a review. *Fundam Clin Pharmacol* 2005;19:27-44.
4. Paul Vanberg, Dan Atar. Androgenic anabolics steroid abuse and the cardiovascular system. *Handb Exp Pharmacol* 2010;411-57.
5. Achar S, Rostamian A, Narayan SM. Cardiac and metabolic effects of anabolic-androgenic steroid abuse on lipids, blood pressure, left ventricular dimensions, and rhythm. *Am J Cardiol* 2010;106:893-901.
6. Nieminen MS, Ramo MP, Viitasalo M, Heikkila P, Karjalainen J, Mantysaari M, et al. Serious cardiovascular side effects of large doses of anabolic steroids in weightlifters. *Eur Heart J* 1996;17:1576-83.
7. D'Andrea A, Caso P, Salerno G, Scarafile R, De Corato G, Mita C, et al. Left ventricular early myocardial dysfunction after chronic misuse of anabolic androgenic steroids: a Doppler myocardial and strain imaging analysis. *Br J Sports Med* 2007;41:149-55.
8. Nottin S, Nguyen LD, Terbah M, Obert P. Cardiovascular effects of androgenic anabolic steroids in male bodybuilders determined by tissue Doppler imaging. *Am J Cardiol* 2006;97:912-5.
9. Krieg A, Scharhag J, Albers T, Kindermann W, Urhausen A. Cardiac tissue Doppler in steroid users. *Int J Sports Med* 2007;28:638-43.
10. Abhayaratna WP, Seward JB, Appleton CP, Douglas PS, Oh JK, Tajik AJ, et al. Left atrial size: physiologic determinants and clinical applications. *J Am Coll Cardiol* 2006;47:2357-63.
11. Sauter HJ, Dodge HT, Johnston RR, Graham TP. The relationship of left atrial pressure and volume in patients with heart disease. *Am Heart J* 1964;67:635-42.
12. Matsuda Y, Toma Y, Ogawa H, Matsuzaki M, Katayama K, Fujii T, et al. Importance of left atrial function in patients with myocardial infarction. *Circulation* 1983;67:566-71.
13. D'Andrea A, De Corato G, Scarafile R, Romano S, Reigler L, Mita C, et al. Left atrial myocardial function in either physiological or pathological left ventricular hypertrophy: A two-dimensional speckle strain study. *Br J Sports Med* 2008;42:696-702.
14. Cameli M, Caputo M, Mondillo S, Ballo P, Palmerini E, Lisi M, et al. Feasibility and reference values of left atrial longitudinal strain imaging by two-dimensional speckle tracking. *Cardiovasc Ultrasound* 2009;8:6.
15. Altekin RE, Yanikoglu A, Karakas MS, Ozel D, Kucuk M, Yilmaz H, et al. Assessment of left atrial dysfunction in obstructive sleep apnea patients with the two dimensional speckle-tracking echocardiography. *Clin Res Cardiol* 2012;101:403-13.
16. Lang RM, Badano LP, Mor-Avi V, Afilalo J, Armstrong A, Ernande L, et al. Recommendations for cardiac chamber quantification by echocardiography in adults: an update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr* 2015;28:1-39.e14.

17. Nagueh SF, Appleton CP, Gillebert TC, Marino PN, Oh JK, Smiseth OA et al. Recommendations for the evaluation of left ventricular diastolic function by echocardiography. *J Am Soc Echocardiogr* 2009;22:107-33.
18. Peterson LR, Waggoner AD, Schechtman KB, Meyer T, Gropler RJ, Barzilai B, et al. Alterations in left ventricular structure and function in young healthy obese women: assessment by echocardiography and tissue Doppler imaging. *J Am Coll Cardiol* 2004;43:1399-404.
19. Kircher B, Abbott JA, Pau S, Gould RG, Himelman RB, Higgins CB, et al. Left atrial volume determination by biplane two dimensional echocardiography: validation by cine computed tomography. *Am Heart J* 1991;121:864-71.
20. Paraskavidis I, Doduras T, Adamopoulos S, Kremastinos DT. Left atrial functional reserve in patients with nonischemic dilated cardiomyopathy: an echocardiographic dobutamine study. *Chest* 2002;122:1340-7.
21. Moyssakis I, Papadopoulos DP, Kelepeshis G, Gialafos E, Votteas V, Triposkiadis F. Left atrial systolic reserve in idiopathic vs. ischemic dilated cardiomyopathy. *Eur J Clin Invest* 2005;35:355-61.
22. Vianna-Pinton R, Moreno CA, Baxter CM, Lee KS, Tsang TS, Appleton CP. Two-dimensional speckle-tracking echocardiography of the left atrium: feasibility and regional contraction and relaxation differences in normal subjects. *J Am Soc Echocardiogr* 2009;22:299-305.
23. Kim DG, Lee KJ, Lee S, Jeong SY, Lee YS, Choi YJ, et al. Feasibility of two-dimensional global longitudinal strain and strain rate imaging for the assessment of left atrial function: a study in subjects with a low probability of cardiovascular disease and normal exercise capacity. *Echocardiography* 2009;26:1179-87.
24. D'Andrea A, Limongelli G, Caso P, Sarubbi B, Della Pietra A, Brancaccio P, et al. Association between left ventricular structure and cardiac performance during effort in two morphological forms of athlete's heart. *Int J Cardiol* 2002;86:177-84.
25. Pelliccia A, Culasso F, Di Paolo F, Maron BJ. Physiologic left ventricular cavity dilation in elite athletes. *Ann Intern Med* 1999;130:23-31.
26. Pluim BM, Zwinderman AH, van der Laarse A, van der Wall EE. The athlete's heart. A metaanalysis of cardiac structure and function. *Circulation* 2000;101:336-42.
27. Richand V, Lafitte S, Reant P, Serri K, Lafitte M, Brette S, et al. An ultrasound speckle tracking (two-dimensional strain) analysis of myocardial deformation in professional soccer players compared with healthy subjects and hypertrophic cardiomyopathy. *Am J Cardiol* 2007;100:128-32.
28. Gabrielli L, Enríquez A, Córdova S, Yáñez F, Godoy I, Corbalán R. Assessment of left atrial function in hypertrophic cardiomyopathy and athlete's heart: a left atrial myocardial deformation study. *Echocardiography* 2012;29:943-9.
29. De Marchi SF, Allemann Y, Seiler C. Relaxation in hypertrophic cardiomyopathy and hypertensive heart disease: Relations between hypertrophy and diastolic function. *Heart* 2000;83:678-84.
30. Scharhag J, Schneider G, Urhausen A, Rochette V, Kramann B, Kindermann W. Athlete's heart: Right and left ventricular mass and function in male endurance athletes and untrained individuals determined by magnetic resonance imaging. *J Am Coll Cardiol* 2002;40:1856-63.
31. Schmidt-Trucksäss A, Schmid A, Häussler C, Huber M, Huonker M, Keul J. Left ventricular wall motion during diastolic filling in endurance-trained athletes. *Med Sci Sports Exerc* 2001;33:189-95.
32. Pearson AC, Schiff M, Mrosek D, Labovitz AJ, Williams GA. Left ventricular diastolic function in weight lifters. *Am J Cardiol* 1986;58:1254-9.
33. Urhausen A, Holpes R, Kindermann W. One- and two-dimensional echocardiography in bodybuilders using anabolic steroids. *Eur J Appl Physiol Occup Physiol* 1989;58:633-40.
34. Urhausen A, Albers T, Kindermann W. Are the cardiac effects of anabolic steroid abuse in strength athletes reversible? *Heart* 2004;90:496-501.
35. Palka P, Lange A, Fleming AD, Donnelly JE, Dutka DP, Starkey IR, et al. Differences in myocardial velocity gradient measured throughout the cardiac cycle in patients with hypertrophic cardiomyopathy, athletes and patients with left ventricular hypertrophy due to hypertension. *J Am Coll Cardiol* 1997;30:760-8.
36. Shan K, Bick RJ, Poindexter BJ, Shimoni S, Letsou GV, Reardon MJ, et al. Relation of tissue Doppler derived myocardial velocities to myocardial structure and beta-adrenergic receptor density in humans. *J Am Coll Cardiol* 2000;36:891-6.
37. To AC, Flamm SD, Marwick TH, Klein AL. Clinical utility of multimodality LA imaging: assessment of size, function, and structure. *JACC Cardiovasc Imag* 2011;4:788-98.
38. Cianciulli TF, Saccheri MC, Lax JA, Bermann AM, Ferreira DE. Two-dimensional speckle tracking echocardiography for the assessment of atrial function. *World J Cardiol* 2010;2:163-70.
39. Saraiva RM, Demirkol S, Buakhamsri A, Greenberg N, Popović ZB, Thomas JD, et al. Left atrial strain measured by two-dimensional speckle tracking represents a new tool to evaluate left atrial function. *J Am Soc Echocardiogr* 2010;23:172-180.
40. Sun P, Wang ZB, Li JX, Nie J, Li Y, He XQ, et al. Evaluation of left atrial function in physiological and pathological left ventricular myocardial hypertrophy by real-time tri-plane strain rate imaging. *Clin Cardiol* 2009;32:676-83.
41. Alizade E, Avcı A, Fidan S, Tabakçı M, Bulut M, Zehir R, et al. The Effect of chronic anabolic-androgenic steroid use on Tp-E interval, Tp-E/Qt Ratio, and Tp-E/Qtc Ratio in Male Bodybuilders. *Ann Noninvasive Electrocardiol* 2015;20:592-600.
42. Ogawa K, Hozumi T, Sugioka K, Iwata S, Otsuka R, Takagi Y, et al. Automated assessment of left atrial function from time-left atrial volume curves using a novel speckle tracking imaging method. *J Am Soc Echocardiogr* 2009;22:63-9.
43. Otani K, Takeuchi M, Kaku K, Haruki N, Yoshitani H, Tamura M, et al. Impact of diastolic dysfunction grade on left atrial mechanics assessed by two-dimensional speckle tracking echocardiography. *J Am Soc Echocardiogr* 2010;23:961-7.
44. Montisci M, El Mazloum R, Cecchetto G, Terranova C, Ferrara SD, Thiene G, et al. Anabolic androgenic steroids abuse and cardiac death in athletes: morphological and toxicological findings in four fatal cases. *Forensic Sci Int* 2012;217:e13-8 (21).
45. Fineschi V. Chronic, supra-physiological doses of nandrolone decanoate and exercise induced cardio-toxicity in an animal-model study. *Acta Physiol (Oxf)* 2013;208:141-3.
46. Montisci R, Cecchetto G, Ruscazio M, Snenghi R, Portale A, Viel G, et al. Early myocardial dysfunction after chronic use of anabolic androgenic steroids combined pulse-wave-tissue doppler imaging and ultrasonic integrated backscatter cyclic variations analysis. *J Am Soc Echocardiogr* 2010;23:516-22.
47. Morris DA, Gailani M, Vaz Pérez A, Blaschke F, Dietz R, Haverkamp W, et al. Left atrial systolic and diastolic dysfunction in heart failure with normal left ventricular ejection fraction. *J Am Soc Echocardiogr* 2011;24:651-62.
48. Kuppahally SS, Akoum N, Burgon NS, Badger TJ, Kholmovski EG, Vijayakumar S, et al. Left atrial strain and strain rate in patients with paroxysmal and persistent atrial fibrillation. Relationship to left atrial structural remodeling detected by delayed-enhancement MRI. *Circ Cardiovasc Imaging* 2010;3:231-9.
49. Ohtani K, Yutani C, Nagata S, Koretsune Y, Hori M, Kamada T. High prevalence of atrial fibrosis in patients with dilated cardiomyopathy. *J Am Coll Cardiol* 1995;25:1162-9.
50. Amundsen BH, Helle-Valle T, Edvardsen T, Torp H, Crosby J, Lyseggen E, et al. Noninvasive myocardial strain measurement by speckle tracking echocardiography: validation against sonomicrometry and tagged magnetic resonance imaging. *J Am Coll Cardiol* 2006;47:789-93.