

PROTECTIVE EFFECTS OF TRIMETAZIDINE IN TRANSIENT SPINAL CORD ISCHEMIA

A. BALTALARLI, MD*,
İ. GÖKŞİN, MD*,
M.H. US, MD**,
G. ÖNEM MD***,
R. ORTAÇ, MD****,
E. COŞKUN, MD*****,
B. H.ŞİRİN, MD*

From:

* Department of
Cardiovascular Surgery,
Pamukkale University,
Faculty of Medicine,
Denizli, Turkey,

** Department of
Cardiovascular Surgery,
GATA Haydarpaşa
Hospital, İstanbul, Turkey,

*** Department of
Pathology, İzmir Behçet Uz
Hospital, İzmir, Turkey,

**** Department of
Cardiovascular Surgery,
İzmir Atatürk State Hospital
İzmir, Turkey,

***** Department of
Neurosurgery, Pamukkale
University, Faculty of
Medicine, Denizli, Türkiye

Address for

reprints:

Dr. Ahmet Baltalarlı
PK. 283 20100
Denizli Türkiye
Tel : +90 258 2131577
Fax: +90 258 2132016
e-mail:
ahmetbaltalarli@superonline.com

The neuroprotective effect of trimetazidine (TMZ) was tested prospectively in a rabbit spinal cord ischemia model. Ischemia was induced by clamping the aorta just distal to the left renal artery and proximal to the aortic bifurcation for a duration of 20 minutes. Twenty five male New Zealand rabbits were randomized as follows: TMZ group (n=10) receiving 3 mg/kg trimetazidine intravenously before the occlusion of aorta, control group undergoing occlusion but receiving no pharmacological intervention (n=10), sham operation group (n=5) subjected to operative dissections without aortic occlusion. Physiological parameters and somatosensory evoked potentials (SEP) were monitored in the animals before and during ischemia and in the 1st, 15th, 60th minutes of reperfusion. Neurologic status was assessed 24 and 48 hours after the operation. Their spinal cord, abdominal aorta, and its branches were processed for histopathologic examination at 48th hours after the operation. At the end of the ischemic period, the average N1-P1 amplitude was reduced to 22% of the baseline in all ischemic animals. This was followed by a gradual return to 90±2% of the initial amplitude in TMZ group and 81±2% in control group (p<0.05) after 60 minutes of reperfusion. The average motor function score was significantly higher in TMZ group than control group (3.7±0.5 vs 3.1±0.6 at 24th and 3.5±0.7 vs 2.9±0.6 at 48th hours, p<0.05). Histologic observations were clearly correlated with neurologic findings. The results suggest that trimetazidine reduces spinal cord injury during thoracoabdominal aortic operations and may have therapeutic utility during high risk operations.

Key words: Trimetazidine, spinal cord ischemia, somatosensory evoked potentials.

Spinal cord ischemia is a devastating complication of thoracoabdominal aortic operations, with paraplegia occurring after as many as 31% of procedures (1). Temporary aortic occlusion may produce a critical reduction in spinal cord perfusion with the risk of irreversible ischemia injury. Reperfusion may also cause numerous further negative effects in case of severe and prolonged ischemia. The molecular and cellular mechanisms that underlie ischemia-reperfusion injury has a multifactorial etiology. During spinal cord ischemia, ischemic cascade starts and results in cellular metabolic dysfunction.

The reasons of reperfusion injury are infiltration of leukocytes, lipid peroxidation, calcium paradox with subsequent release of free oxygen radicals into the previously ischemic tissue (2,3). Pretreatment with some pharmacological agents to reduce or eliminate ischemic injury could be of great benefit in cardiovascular surgery.

Trimetazidine (TMZ), 1-(2,3,4 trimethoxybenzyl) piperazine dihydrochloride, an antianginal drug used in clinical treatment of ischemia, has been shown to act at different levels of sequence of events that occur during ischemia. It improves the efficiency of oxygen consumption and protects cardiac cells against the fall in intracellular ATP (4), reduces the release of phosphocreatine (5) and has an antioxidant activity (6,7).

The aim of this study was to determine whether trimetazidine administered before aortic occlusion was capable of preserving neurological status and maintaining the structural integrity of the spinal cord in severe ischemia (8).

MATERIALS AND METHOD

Animal care and surgical technique:

Our experimental protocol was approved by the Animal Care and Use Committee of Pamukkale University. Experiments were performed on 25 male New Zealand white rabbits weighing 3 to 4 kg. The animals were housed in individual cages under controlled conditions (12 hourly light and dark cycle and room temperature of 21°C) and free access to water and laboratory chow. During the surgical procedures, anesthesia was induced with intraperitoneal ketamine (50 mg/kg) and xylazine (10 mg/kg), followed fractionally by the cocktail of ketamine (16 mg/kg) and xylazine (3 mg/kg). No animals received hemodynamic or ventilatory support. The animals were placed in a nose cone to breathe oxygen at a rate of 2L/minute. Rectal temperature was monitored and maintained close to 38°C under a warming light. An intravenous catheter (24 gauge) was placed in an ear vein, and preoperatively cefazolin 50 mg/kg was given as a single dose. Maintenance fluid of 0.9% NaCl was infused at a rate of 20 ml/h during the procedure. The

right femoral artery was exposed for arterial line with a catheter and connected to blood pressure/heart transducer and monitored (Hewlett-Packard monitor, model 78352C, Germany). Percutaneous arterial oxygen saturation was continuously monitored.

Experimental groups and surgical procedures:

Rabbits were randomly allocated into three groups. In TMZ group (n=10), trimetazidine (Servier, France) was injected as a bolus (3 mg/kg) via ear vein immediately after anesthesia. Control animals (n=10) underwent standard aortic occlusion and intravenous injection of 0.9% sodium chloride under identical conditions to that of the trimetazidine injection group.

After the surgical preparation using aseptic techniques, a midline laparotomy was performed, and then infrarenal aorta and aortic bifurcation were exposed. Intravenous heparin (150 U/kg) was given and spinal cord ischemia was induced by cross-clamping of the aorta with two bulldog clamps for 20 minutes. The second clamp was used for occluding iliac collateral circulation. After the ischemic period, the clamps were removed and restoration of blood was verified visually. The abdomen was then closed. In sham operation group, incision was left open for 20 minutes corresponding to spinal cord ischemia period but aorta was not occluded. This group of animals was used for observing the effects of anesthesia and operation on results.

SEP recording:

In order to assess neurologic recovery, somatosensory evoked potentials (SEP) were monitored before the occlusion of the abdominal aorta, during ischemia and at the first, 10th and 60th minutes of reperfusion. SEP responses were evaluated for re-establishment of the percentages of their baseline control values. Medelec Premiere Plus (Medelec Ltd., UK) was used for the recordings.

The recording electrodes were inserted percutaneously just lateral to the spinous process of T2 vertebra, so that the tips rested on the lamina. To record the SEPs, the left

sciatic nerve was stimulated by a bipolar electrode with square wave pulses of 0.1 msec duration and approximately 7 mA intensity delivered at 3 Hz. The filter range was 20 Hz to 1 kHz. Sixty four responses were averaged and displayed. The latencies and amplitudes of SEP before and after the ischemic procedure were compared.

Postoperative care and assessment:

Femoral arterial and venous lines were removed at the 60th minute of reperfusion. When the animals awakened from anesthesia, they were returned to their cages. The Crede maneuver was used to empty the bladders of paraplegic animals at least twice daily.

Neurologic status was scored by assessment of hind limb neurologic function according to the Tarlov scale (9) [0: no movement; 1: slight movement; 2: sits with assistance; 3: sits alone; 4: weak hop; 5: normal hop] at the 24th and 48th hours after ischemia. Two observers (1 of whom was blinded to the experimental conditions) graded the neurologic status independently.

Histopathology:

After the last neurological examination at the 48th hour, the animals were anesthetized and killed with an intracardiac perfusion of 10% neutral formalin. The retroperitoneal region including the abdominal aorta and its branches were removed, extensively and fixed in 10% formalin solution. The spinal cord was then removed, immersed in the same fixative, and postfixed for about 14 days before being set in paraffin blocks for sectioning. Section of the thoracic and lumbar cord were stained with hematoxylin-eosin. The abdominal aorta and its branches were examined for revealing possible thrombosis or embolic occlusion.

Statistical Analysis:

Statistical analysis of physiologic and hemodynamic parameters were performed by analysis of variance for repeated measures. All parametric data were expressed as mean \pm standard deviation. Statistical analyses of the neurologic

scores were done with the non-parametric Man-Whitney U test using the Bonferroni correction. Somatosensory evoked potential data are expressed as percent of baseline control value. Recovery of SEP amplitude in trimetazidine treated animals was compared with that of ischemia control animals using t-test and Bonferroni correction. A p value of less than 0.05 was considered significant

RESULTS

No differences in mean levels of blood glucose, arterial blood gases, pH, rectal temperature, heart rate, and mean arterial blood pressure were noted among groups. The femoral artery blood pressures were measured near zero in the animals subjected to ischemia during the ischemic period and became gradually normal within the first minute of beginning of reperfusion and corrected with intravenous sodium bicarbonate (1.7 ± 0.4 mEq in TMZ group and 1.8 ± 0.4 mEq in control group, statistically nonsignificant) and no correction was required in the sham group.

SEP recording showed a negative peak (N1) with an average latency of 5.3 ± 0.2 msec, a positive peak (P1) with an average latency of 6.3 ± 0.1 msec and a negative peak (N2) with average latency of 10.1 ± 0.2 msec. N1-P1 amplitude was $5.2 \pm 0.4 \mu V$ and P1-N2 amplitude was $7.4 \pm 0.3 \mu V$. The peaks were stable in sham group during the procedure. In the ischemic groups, N1 and P1 peaks progressively declined and with an average of 8.1 \pm 4 minute after aortic occlusion. At the end of the ischemic period, average N1-P1 amplitude decreased 22% pre-ischemic baseline level in all ischemic animals. The average amplitude of P1-N2 dropped to 15% of pre-ischemic baseline level in all ischemic animals. This was followed by a gradual return to $90 \pm 2\%$ of the initial N1-P1 amplitude and $80 \pm 2\%$ P1-N2 amplitude after the 60th minute of reperfusion in the TMZ group. However this recovery were $81 \pm 2\%$ and $73 \pm 2\%$, respectively, of the initial amplitude in the control group. There was a corresponding statistical difference between the TMZ and the control group ($p > 0.05$). The




Figure 1. Typical SEP traces of an ischemic animal.

A: Pre-ischemic baseline SEP trace, and N1-P1 and P1-N2 amplitudes,
 B: SEP traces during ischemia,
 C: SEP trace at the 1st minute of reperfusion,
 D: SEP trace at the 15th minute of reperfusion,
 E: SEP trace at the 60th minute of reperfusion

peaks were stable in the sham operated group during the procedure. SEP recordings before and after the occlusion of the abdominal aorta and in the reperfusion period are illustrated in Figure 1.

Neurologic outcomes:

Sham operated animals showed full neurological recovery after the procedure. The evaluation of motor function score at the 24th and 48th hours of post-ischemic period in the TMZ and control groups is presented in Table 1. All ischemic animals exhibited neurologic

deficiencies. The average motor score of the TMZ group was significantly ($p < 0.05$) better than the control group 24 and 48 hours after the ischemia.

Histopathological findings:

Histopathologic examination of abdominal aorta and its branches were normal and revealed no thrombus formation in all animals. The histopathologic findings of spinal cord correlated with the neurologic outcomes. The animals with high motor score exhibited unaltered structure of gray matter or minimal necrosis (Figure 2). But the animals with low motor score had typical infarctions of variable size in the central gray matter (Figure 3).




Figure 2. Photomicrograph of the cells in the anterior horn of spinal cord showing karyolysis and moderate changes in the cytoplasm (TMZ group, score 4 of motor neurological function, lumbosacral region, X200 hematoxylin-eosin).

Table 1. Tarlov scores of the TMZ and control groups at 24th and 48th hours after the ischemia.

Animal groups	n	Motor score					Average motor score
		0	1	2	3	4	
TMZ group							
24 th h	10				3	7	3.7±0.5*
48 th h	10			1	3	6	3.5±0.7*
Control group							
24 th h	10			1	7	2	3.1±0.6
48 th h	10			2	7	1	2.9±0.6

* Significantly different from control group ($p < 0.05$) by Mann-Whitney U test.




Figure 3. Neuronal death characterized by blurred cytoplasmic border, degranulation of Nissl bodies, karyolysis, pericellular vacuolization (control group, score 2 of motor neurological function, lumbosacral segment, X400 hematoxylin-eosin).

DISCUSSION

Numerous surgical techniques and pharmacological interventions have been used to reduce the rate of perioperative paraplegia after operations on the thoracoabdominal aorta, but none can claim consistently proven efficacy (10). The data presented in this study demonstrated that reduction in neuronal damage has been achieved with pre-ischemic, intravenous administration of trimetazidine in a rabbit model of transient global ischemia. Animals that were subjected to 20 min of ischemia after receiving 3 mg/kg trimetazidine exhibited better neurologic outcomes. There are very little knowledge in the literature about neuroprotective effect of trimetazidine, although there is a great deal of studies about its cardioprotective effect. The efficacy of trimetazidine has recently been investigated on reducing organ damage in myocardium (5) but not in spinal cord ischemia.

In the presence of cellular ischemia, there is a deterioration of energy production, distortions of cellular homeostasis (acidosis and cytoplasmic sodium and calcium accumulation) and an overproduction of free radicals which have detrimental effects on cell structure and function. Trimetazidine antagonizes hypoxia induced reduction in

myocardial ATP levels without diminishing phosphocreatine levels (11), and accelerates the reconstitution of energy pools during reperfusion perhaps by inhibiting mitochondrial ATPase (12). Under normal physiological conditions, trimetazidine does not affect the plasma membrane ionic channels or ionic pumps in the cardiac cells. But under ischemic conditions it reduces myocardial intracellular acidosis, and indirectly limits the accumulation of Na^+ and Ca^{2+} (13, 14).

Another mechanism is that trimetazidine significantly reduces H_2O_2 production (13,15) and finally it decreases adhesiveness and aggregation of platelets which may be due to a blockade of the process preceding the cyclooxygenase pathway and related to the free-radical scavenging property of this compound (16, 17).

According to Harpey et al. (15), there is no major difference in pharmacokinetic parameters between single or repeated doses. So, we decided to administer trimetazidine as single bolus treatment. Trimetazidine at intravenous doses from 0.25 to 3.0 mg/kg, do not modify hemodynamic parameters in the systemic or coronary circulation (13,15). We also had similar hemodynamic findings on the systemic circulation with a dose of 3 mg/kg. Although oral administration of trimetazidine is widely used in clinical practice, there are some literature on the effectiveness of parenteral route and bolus dose (10,18-21).

The present study was similar to the clinical setting of thoracoabdominal aortic surgery leading to spinal cord ischemia. There was no observed significant difference in blood pressure and blood glucose levels among all comparable groups. This model might increase the tolerance of spinal cord to ischemia before elective surgery.

The rabbit spinal cord is a reliable model for systematically and rapidly observing the protective effects of investigated agents on ischemia and reperfusion injury. However, ligation of the rabbit's abdominal aorta may not produce complete ischemia in the spinal cord. There are considerable individual variations in residual collateral blood flow, and in most animals, about 2% of normal blood flow remains after 20 minutes of ligation (9). In the present study, we used a

second clamp at the distal abdominal aorta for occluding the iliac collateral circulation and monitored the spinal cord ischemia with SEP. Monitoring of SEP during aortic occlusion has been used for identification and exclusion of animals which have sufficient collateral blood flow to permit the spinal cord to survive long periods of aortic occlusion (22). Previous studies have shown that a decrease of 15-20% in amplitude can be taken as a criteria of significant SEP change (23). In the present study, despite minimal individual variations, SEP amplitudes decreased to about 22% of their pre-ischemic baseline values in most of the animals at 20th min of ischemia, and monitoring of SEP in this model has permitted observation of similar collateral blood flow properties in the studied groups of animals. In conclusion, this study indicated that trimetazidine potentially ameliorated reperfusion injury in a rabbit model of transient spinal cord ischemia, and animals treated with trimetazidine showed better neurologic outcome. In case of neurologic deficit caused by aortic dissection, this drug may be used until reperfusion is able to be established. In addition, trimetazidine can be used in the prevention of spinal cord injury during thoracoabdominal aortic operations.

REFERENCES

1. Svenson LG, Crawford ES, Hess KR, Coselli JS, Safi HJ. Experience with 1509 patients undergoing thoracoabdominal aortic operations. *J Vasc Surg* 1993;17:350-70.
2. Harlan JM. Leucocyte-endothelial interactions. *Blood* 65:513-25.
3. Matsuo Y, Onodera H, Shiga Y, Nakamura M, Ninomiya M, Kihara T, Kogure K. Correlation between myeloperoxidase-quantified neutrophil accumulation and ischemic brain injury in the rat. *Stroke* 1985;25:1469-75.
4. Lavancy N, Martin J, Rossi A. Anti-ischemic effects of trimetazidine 31 P-NMR Spectroscopy in the isolated rat heart. *Arch Int Pharmacodyn Ther* 1987;286:97-110.
5. Libersa C, Honore E, Adamantidis M, Rouet E, Dupuis B. Antiischemic: effect of trimetazidine enzymatic and electric response in a model of in-vitro myocardial ischemia. *Cardiovasc Drug Rev* 1990;4:808-9.
6. Harpey C, Maridonneau-Parini I. Trimetazidine protects the human red blood cell against oxygen free radical damage. *Cardiovasc Drug Rev* 1990;4:818-9.
7. Renaud JF. Internal pH, Na, and Ca regulation by trimetazidine during cardiac cell acidosis. *Cardiovasc Drug Rev* 1988;4:677-85.
8. DeGirolami U, Zivin JA. Spinal cord infarction: A highly reproducible stroke model. *Stroke* 1980;11:200-2.
9. Tarlov IM. Acute spinal cord compression on paralysis. *J Neurosurg* 1972;36:10-20.
10. Gharagozloo F, Larson J, Dausmann MJ, Neville RF Jr, Gomes MN. Spinal cord protection during surgical procedures on descending thoracic and thoracoabdominal aorta: review of current techniques. *Chest* 1996;109:799-809.
11. Rochette L, Fitoussi M, Bralet J. Effects du pre-traitement par la trimetazidine sur le metabolisme energetique du coeur isole de rat soumis a une ligature coronaire et perfuse en normoxie ou en hypoxie. *Gaz Med Fr* 1984;91(suppl):17-21.
12. Cruz C, Zaoui A, Ayoub S, Harpey C, Goupt P, Younes A. Alterations des myocytes isoles des ventricules de coeur de rat adulte: protection par la trimetazidine. *Concours Med* 1987;36 (suppl):3470-5.
13. Clauser P, Harpey C. Antioxidant properties of an anti-ischemic agent: trimetazidine. *Adv Exp Med Biol* 1990;264:247-50.
14. Yanagisawa, Satoh K, Taira N. Simultaneous assessment of the effects of trimetazidine on myocardium and coronary vasculature of the dog. *Arch Int Pharmacodyn Ther* 1979;237:316-28.
15. Harpey C, Clauser P, Labrid C, Freyria JL, Poirier JP. Trimetazidine, a cellular anti-ischemic agent. *Cardiovasc Drug Rev* 1989;6:292-312.

16. Higuchi H, Hayashi M, Sano I: Effect of trimetazidine granules (Vastarel 20 mg) in platelet adhesiveness and aggregation in patients with ischemic heart disease. Response to exercise. *J Nissei Hosp* 1981;9:89-95.
17. Shirahase H, Suzuki Y, Osum S, Kakaya N, Kurahashi K: Inhibitory effects of trimetazidine dihydrochloride on aggregation, serotonin release and malodialdehyde production in rabbit platelets. *Jpn J Pharmacol* 1988;47:29-34.
18. Kober G, Pennaforte S, Buck T, Sievert H, Vallbracht C: Myocardial cytoprotection during percutaneous transluminal coronary angioplasty. *Eur Heart J* 1992;13:1109-15.
19. Özden A, Aybek Z, Saydam N, Çallı N, Saydam O, Düzcan E, Güner G: Cytoprotective effect of trimetazidine on 75 min warm renal ischemia-reperfusion injury in rats. *Eur Surg Res* 1998;30:227-34.
20. Sakai K, Fukushi Y, Abiko Y: Inhibitory effect of trimetazidine on utilization of myocardial glycogen during coronary ligation in dogs. *Pharmacology* 1986;32:72-93.
21. Williams FM, Tanda K, Kus M, Williams TJ: Trimetazidine inhibits neutrophil accumulation after myocardial ischemia and reperfusion in rabbits. *J Cardiovasc Pharmacol* 1993;22:828-33.
22. Robertson CS, Foltz R, Grossman RG, Goodman JC: Protection against experimental ischemic spinal cord injury. *J Neurosurg* 1986;46:633-42.
23. Stechison MT, Panagis SG, Reinhart SS: Somatosensory evoked potential monitoring during spinal surgery. *Acta Neurochir (Wien)* 1995;135:56-61.