

## SS-03

### Percutaneous Coronary Intervention with the DK-Crush Technique for a Complex LAD-Diagonal Bifurcation Lesion in the Setting of Acute Anterior Myocardial Infarction

Ceyda Nur Batak, Fatih Kahraman

Kütahya Health Sciences University, Kütahya, Türkiye

**Introduction and Aim:** Coronary bifurcation lesions account for approximately 15–20% of all percutaneous coronary interventions (PCI). True bifurcation lesions of the left anterior descending artery (LAD) and a large diagonal branch are among the most technically demanding PCI scenarios in acute ST-elevation myocardial infarction (STEMI). Double kissing (DK) crush has been associated with superior outcomes compared with other two-stent techniques in complex bifurcation lesions; however, repeated wire crossings and kissing balloon steps may create technical difficulties in the time-constrained acute setting. We present an LAD-second diagonal (D2) true bifurcation lesion treated with DK-crush during primary PCI, complicated by three distinct technical challenges.

**Case Report:** A 66-year-old man with no known cardiac history presented with chest pain and ST elevation in the anterior leads on electrocardiography. Blood pressure was 141/93 mmHg, heart rate 78 bpm, oxygen saturation 97%. Echocardiography showed an ejection fraction of 40% with anterior wall and apical hypokinesia. Emergency angiography via a right femoral 7F sheath revealed a markedly elongated aorta and a superiorly originating left main coronary artery; a true bifurcation lesion with 90% calcific stenosis in the LAD proximal to the D2 origin and 90% stenosis in the proximal D2 was identified, and DK-crush was planned. Both branches were wired separately through an EBU 4.5 guiding catheter. During predilatation, refractory loss of flow developed in D2; as repeated prolonged balloon dilatations failed, 10 mg of intracoronary tissue plasminogen activator was administered within 10 minutes, and flow was restored after prolonged inflation with a 2.75×30 mm non-compliant (NC) balloon. A 2.75×22 mm drug-eluting stent was implanted in D2, followed by proximal optimization technique (POT) and the first kissing balloon. After the first crush, a 3.0×32 mm stent could not be advanced into the LAD; after exchange for a more supportive wire with microcatheter support and re-crushing of the side branch stent with a 4.0×8 mm NC balloon, the second kissing balloon was performed and the stent implanted successfully. Repeat POT and kissing balloon provided complete angiographic patency of the bifurcation. At the end of the procedure, 99% ostial stenosis (jailing) developed in a well-developed first diagonal branch (D1) arising proximal to the stent; this branch was rewired and opened with 2.0×10 mm balloon angioplasty without stent deformation. TIMI 3 flow was obtained in the LAD, D1 and D2. He was monitored in the coronary intensive care unit under inotropic support; apart from a self-terminating 10–15-second episode of non-sustained ventricular tachycardia, no complications occurred. He was discharged on acetylsalicylic acid, prasugrel, rosuvastatin, ramipril and metoprolol, with outpatient follow-up at 1 month.

**Conclusion:** LAD-diagonal true bifurcation lesions can be treated successfully with DK-crush in acute STEMI; however, difficult stent delivery in calcified and tortuous anatomy, refractory side branch no-reflow and jailing of an adjacent side branch should be anticipated. This case suggests that low-dose intracoronary thrombolysis added to prolonged balloon dilatation in refractory no-reflow, and wire exchange with a repeat crush maneuver when the stent cannot be advanced after the first crush, may be effective bailout strategies.

## SS-04

### Dual Plaque Modification Strategy in High-Risk Complex PCI

Eda Özcan, Beliz Koçyiğit, Hakan Erkan

Bursa City Hospital, Bursa, Türkiye

**Introduction and Aim:** The success of percutaneous interventions in fragile patients with severely calcified and multivessel coronary artery disease depends on optimized lesion preparation. When the calcific burden is extremely high, a single device may not be sufficient. In this case report, we aim to present a complex percutaneous coronary intervention (PCI) procedure successfully managed by combining two different advanced plaque modification methods—rotablation and a cutting balloon—in the same session in a patient with active malignancy and end-stage renal failure who refused surgery.

**Case Report:** A 79-year-old female patient presented to the emergency department with shortness of breath. Her medical history included diabetes mellitus, hypertension, chronic kidney disease (stage 3/7 hemodialysis), and a history of breast cancer (right mastectomy in 2024). The patient, whose ECG showed normal sinus rhythm with no ST-segment elevation, was admitted with a diagnosis of non-ST-elevation acute coronary syndrome (NSTEMI-AKS). Transthoracic echocardiography showed normal left ventricular (LV) dimensions and systolic functions (ejection fraction of 55–60%) along with normal right-sided structures. While degenerative changes were observed in the valves, a gradient of 15/26 mmHg on the basis of calcific aortic degeneration, mild aortic regurgitation, mild-to-moderate mitral regurgitation, and mild tricuspid regurgitation were detected. Thorax CT revealed bilateral pleural effusion and cardiomegaly. Coronary angiography (CAG) revealed a severely calcified and thrombotic long lesion causing 50–60% stenosis in the left main coronary artery (LMCA), 60–70% stenosis in the left anterior descending (LAD) artery, and 80–90% stenosis in the circumflex (CX) artery. Coronary artery bypass grafting (CABG) was recommended by the Heart Team (Consensus); however, the patient and her relatives declined the surgical risk and provided informed refusal for CABG. Consequently, a high-risk complex PCI strategy was decided for the patient. To overcome the heavy calcific lesions, a dual plaque modification strategy was planned. First, rotational atherectomy (rotablation) was performed on the heavy calcific lesion in the LAD using a 2.0 mm burr. Following atherectomy, non-compliant (NC) balloon dilatations were performed, a 2.5×38 mm drug-eluting stent (DES) was implanted distally and a 3.0×28 mm DES proximally, and optimal luminal expansion was achieved by post-dilatations with intra-stent NC balloons. Subsequently, the procedure moved to the CX lesion; predilatation was performed on this lesion sequentially using 1.25, 1.5, 2.0, and 2.5 mm chronic total occlusion (CTO) balloons, followed by 2.5 and 3.0 mm NC balloons. To complete the calcific lesion modification, the second method—the cutting balloon strategy—was initiated. Dilatation was performed using a 4.0×15 mm cutting balloon. Following successful plaque modification, a 4.0×38 mm DES was successfully implanted in the CX artery, and high-pressure dilatation was performed with a 4.5 mm intrastent NC balloon. Complete luminal patency (TIMI-3 flow) was achieved at the end of the procedure, and the patient was discharged stably with a plan for medical follow-up.

**Conclusion:** In patients with heavily calcified multivessel disease, the sequential or combined use of plaque modification methods with different mechanisms, such as rotablation and cutting balloons, is life-saving for stent success and expansion.

**Keywords:** Rotational atherectomy, coronary artery calcification, multivessel coronary disease.

## SS-05

### A Case of Inferior Myocardial Infarction Due to Catheter-Induced Dissection During RCA FFR Assessment

Emrah Kaya, Emre Berk Erkip

Department of Cardiology, Kütahya City Hospital, Kütahya, Türkiye

**Introduction and Aim:** Fractional flow reserve (FFR) is a widely used and reliable method for assessing the functional significance of coronary artery stenoses. However, rare but life-threatening complications such as coronary dissection, spasm, or acute thrombotic occlusion may occur

during the procedure. In this case, we present a catheter-induced dissection of the right coronary artery (RCA) that occurred during FFR assessment in the setting of coronary angiography, followed by acute thrombotic total occlusion, and its management with percutaneous coronary intervention, temporary pacemaker support, and thrombolytic therapy.

**Case Report:** A 54-year-old female patient presented to the cardiology outpatient clinic with exertional dyspnea. Coronary computed tomography revealed coronary artery disease, and she was hospitalized for coronary angiography. Angiography demonstrated 70-80% stenosis in the proximal RCA. The decision was made to assess the severity of the RCA lesion with FFR (Fig. 1). After the FFR pressure wire was advanced distal to the lesion, the patient developed sudden ST-segment elevation and complete atrioventricular (AV) block. Imaging revealed acute total occlusion of the RCA due to a dissection originating from the proximal segment, with a heavy thrombus burden, and the patient sustained an inferior myocardial infarction during the procedure (Fig. 2). The lesion was crossed with a guidewire, and repeated predilations were performed with a 3.0×30 mm balloon. Because of complete AV block, a temporary pacemaker was placed via the right femoral vein. Intracoronary adrenaline was administered for no-reflow. A 4.0×38 mm drug-eluting stent was implanted in the RCA. A second 4.5×12 mm drug-eluting stent was placed at the RCA ostium, overlapping with the first stent. TIMI 2-3 flow was achieved after proximal optimization, and a 12-hour glycoprotein IIb/IIIa inhibitor infusion was initiated (Fig. 3). The patient's rhythm converted from complete block to normal sinus rhythm. On the same day, the patient's general condition deteriorated during follow-up, and complete AV block recurred, prompting a second angiographic evaluation. TIMI 1 flow and dense thrombus were observed in the RCA. Intracoronary alteplase 10 mg was administered over 10 minutes. Although TIMI 3 flow was subsequently achieved, findings consistent with residual thrombus or dissection extending from the mid segment to the distal vessel persisted. The patient then received a 25 mg intravenous alteplase infusion over 24 hours. A third coronary angiography performed two days later showed complete resolution of the thrombosed dissection area distal to the stent, with preserved distal TIMI 3 flow (Fig. 4). The proximal in-stent segment was optimized by high-pressure dilation with a 4.5×15 mm non-compliant balloon. During follow-up, the patient remained in sinus rhythm and hemodynamically stable, and she was discharged on medical therapy.

**Conclusion:** Acute thrombotic total occlusion secondary to dissection occurring during FFR assessment of the RCA is a rare but life-threatening complication that can result in inferior myocardial infarction, complete atrioventricular block, and severe flow impairment. Prompt recognition of electrocardiographic changes and conduction disturbances during the procedure, together with immediate angiographic evaluation, temporary pacemaker support, and timely percutaneous coronary intervention, is of vital importance. In selected cases with refractory and persistent thrombus burden, intracoronary and intravenous alteplase therapy may help restore normal coronary flow while avoiding repeat stent implantation.


**Keywords:** fractional flow reserve, inferior myocardial infarction, coronary artery dissection, intracoronary thrombolytic therapy.



**Figure 1.** RCA lesion before FFR.



**Figure 2.** Totally thrombosed RCA lesion due to dissection caused by the FFR wire.



**Figure 3.** RCA after stent implantation and Gp IIb/IIIa infusion.



**Figure 4.** Control angiography after alteplase infusion, showing resolution of the thrombosed-dissected segment with TIMI 3 flow in the RCA.

## SS-06

### Successful Percutaneous Intervention for a Critical Ostial RCA Lesion Using the Distal Anchor Balloon Technique

Emrah Kaya, Fatih Kahraman

Department of Cardiology, Kütahya Health Sciences University Faculty of Medicine, Kütahya, Türkiye

**Introduction and Aim:** Coronary artery ostial lesions pose significant technical challenges during percutaneous coronary interventions because of inadequate guide catheter support and difficulty advancing devices across the lesion. In such cases, support-enhancing strategies may